

The background of the cover is a photograph of a mountain range. The mountains are rugged and covered in patches of snow, set against a clear blue sky. In the foreground, there is a dense forest of evergreen trees, and below them, a field of low-lying, greyish-green vegetation.

ECOLOGY of NORTH AMERICA

SECOND EDITION

Brian R. Chapman and Eric G. Bolen

WILEY Blackwell

Ecology of North America

*Dedicated to our wives, Sandy and Elizabeth,
whose love, forbearance, and creativeness
inspire and sustain us.*

Ecology of North America

Brian R. Chapman

Senior Research Scientist
Texas Research Institute for Environmental Studies
Sam Houston State University
Huntsville, Texas

Eric G. Bolen

Professor Emeritus
Department of Biology and Marine Biology
University of North Carolina Wilmington
Wilmington, North Carolina

SECOND EDITION

WILEY Blackwell

This edition first published 2015 © 2015 by John Wiley & Sons, Ltd.

Registered Office

John Wiley & Sons, Ltd., The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book.

Limit of Liability/Disclaimer of Warranty: While the publisher and author(s) have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. It is sold on the understanding that the publisher is not engaged in rendering professional services and neither the publisher nor the author shall be liable for damages arising herefrom. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication data

Chapman, Brian R., author.

Ecology of North America / Brian R. Chapman and Eric G. Bolen. – Second edition.

pages cm

Includes bibliographical references and index.

ISBN 978-1-118-97154-3 (cloth)

1. Ecology—North America. I. Bolen, Eric G., author. II. Title.

QH102.B65 2015

577.097—dc23

2015006625

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover image: The spectacular Teton Range in Wyoming offers a centerpiece for the unspoiled places remaining in North America. Lacking foothills, these mountains rise majestically 2200 m (7000 ft) above Jackson Hole that together form Grand Teton National Park—a sanctuary protecting more than 100 species of vascular plants, 300 species of birds, and 60 species of mammals. Some 100 species of grasses and wildflowers enrich the sagebrush flats on the valley floor. Photograph reproduced by permission of Sandra S. Chapman.

Set in 8.5/10.5pt Meridien by SPi Global, Pondicherry, India

Contents

- Foreword, viii
- Preface, ix
- Acknowledgments, xi
- 1 Introduction, 1**
 - A brief overview of ecology, 1
 - The ecosystem, 2
 - Abiotic limits, 3
 - Climate and topography, 4
 - Soils and soil profiles, 4
 - Biotic community, 6
 - Community succession, 6
 - Plant succession: from pioneer to climax, 6
 - Primary and secondary succession, 6
 - Succession and species abundance, 7
 - The biome concept, 7
 - Biodiversity, 7
 - The “species richness gradient”, 8
 - Biodiversity “hotspots”, 8
 - Patterns of distribution, 9
 - Continental patterns, 9
 - Geographical and ecological distribution, 10
 - Some ecological concepts, 10
 - Niches, 10
 - Ecological equivalents, 12
 - Bergmann’s rule, 12
 - Allen’s rule, 13
 - Readings and references, 14
- 2 Tundra, 17**
 - Climatic and glacial influences, 17
 - Soils and geological influences, 17
 - Permafrost, 18
 - Patterned ground, 18
 - Eskers and tundra wildlife, 20
 - Glacial refugia, 21
 - Features and adaptations, 21
 - Plant adaptations to harsh conditions, 21
 - Plant growth and reproduction, 22
 - Decomposition and soil nutrients, 23
 - Some animal adaptations, 23
 - Major vegetative communities, 24
 - Shrub tundra, 25
 - Dwarf birch heath, 25
 - Cottongrass heath, 25
 - Fellfields, 25
 - Invertebrates and tundra ecology, 25
 - Selected tundra mammals, 26
 - Lemmings, 26
 - Arctic ground squirrels, 27
 - Arctic foxes, 28
 - Barren-ground caribou, 28
 - Selected tundra birds, 29
 - Gyrfalcons, 29
 - Snowy owl, 30
 - Ross goose, 30
 - Highlights, 31
 - Absentees: amphibians and reptiles, 31
 - Lichens and “reindeer moss”, 31
 - Snow goose “eat outs”, 32
 - Alpine Tundra, 33
 - Fragile Tundra, 34
 - Impacts of human activity, 34
 - Global warming, 35
 - Readings and references, 36
- 3 Boreal Forest, 41**
 - Climatic boundaries and soils, 41
 - Features and adaptations, 42
 - Plant adaptations, 42
 - Animal adaptations, 42
 - Frequent fires, 42
 - Niches in the Boreal Forest, 43
 - Selected biotic communities, 44
 - Tree line and forest tundra, 44
 - Muskeg, 46
 - Coniferous swamps, 46
 - Comparative ecology of lakes, 46
 - Appalachian Extension, 48
 - Mountain balds, 48
 - Highlights, 49
 - The 10-year cycle, 49
 - Wolves and moose, 50
 - A wealth of salamanders, 52
 - Red squirrels, 53
 - Ecological challenges, 54
 - Acid rain, 54
 - Spruce budworm and DDT, 55
 - Balsam woolly adelgid, 55
 - The Boreal Forest Agreement, 56
 - Readings and references, 57
- 4 Eastern Deciduous Forest, 61**
 - Climatic boundaries and soils, 61
 - Features and adaptations, 61
 - The forest primeval, 63
 - The layered forest, 63

- Autumn leaves, 64
- Ground and leaf litter, 64
- Mast, 65
- Biotic associations, 66
 - Northern hardwoods conifer forests, 66
 - Beech–Maple–Basswood, 67
 - Mesophytic forest, 68
 - Oak–Hickory, 69
 - Mississippi alluvial plain, 70
 - Southern Mixed Forest, 70
- Some associated communities, 71
 - Longleaf pine forests, 71
 - New Jersey Pine Barrens, 73
 - Carolina bays, 74
- Highlights, 75
 - Acorns and blue jays, 75
 - Deer yards, 75
 - Kirtland’s warblers and fire, 76
 - Franklin’s lost tree, 77
 - Cicadas: buzz in the forest, 77
- Ecological challenges, 78
 - Declines of neotropical migrants, 78
 - Forest destruction by exotic organisms, 80
 - Reintroduction of red wolves, 81
- Readings and references, 82
- 5 Grasslands: Plains and Prairies, 89**
 - Major associations, 89
 - Tallgrass prairie, 89
 - Midgrass prairie, 90
 - Shortgrass prairie, 90
 - Transition zones, 93
 - Aspen parklands, 93
 - Cross Timbers, 96
 - Western transition, 96
 - Features and adaptations, 97
 - Seasonal grasses, 97
 - Soils, 99
 - Role of fire, 101
 - Prairie streams, 101
 - Prairie wetlands and waterfowl, 102
 - Pleistocene extinctions, 103
 - Selected prairie mammals, 104
 - Bison, 104
 - Prairie dogs, 104
 - Pronghorns, 106
 - Selected prairie birds, 106
 - Burrowing owls, 106
 - Prairie chickens, 107
 - Highlights, 108
 - Riparian forests, 108
 - The Platte River, 108
 - Nebraska Sandhills, 109
 - Ants, 109
 - Isolation and contact on the plains, 110
 - Grassland settlement, 110
 - Prairie preservation, 111
 - Readings and references, 113
- 6 Regional Grasslands and Related Areas, 120**
 - Regional associations, 120
 - Palouse prairie, 120
 - California Annual Grasslands, 122
 - Southwestern desert grasslands, 124
 - Edwards Plateau, 126
 - Tamaulipan Mezquital, 127
 - Highlights, 129
 - Rodents and vegetation, 129
 - Channeled Scablands, 130
 - Snake River Birds of Prey Conservation Area, 130
 - Mima mounds, 131
 - Desertification, 132
 - Readings and references, 132
- 7 Deserts, 136**
 - Physical geography, 136
 - Why deserts are dry, 136
 - Desert mountains and bajadas, 138
 - Ancient lakes, 138
 - Features and adaptations, 139
 - Desert soils and surfaces, 139
 - Plant adaptations, 140
 - Animal adaptations, 141
 - The major deserts, 144
 - Chihuahuan Desert, 144
 - Sonoran Desert, 145
 - Mojave Desert, 147
 - Great Basin Desert, 149
 - Highlights, 150
 - Nurse trees, 150
 - “Trees” for desert woodpeckers, 151
 - Boojums and elephants: unique trees, 153
 - Yucca moths, 154
 - Desert fishes, 154
 - Realm of reptiles, 155
 - Of soils and mice, 156
 - Deserts and predators, 157
 - Pygmies of the sagebrush steppe, 158
 - Desert quail, rainfall, and vitamin A, 159
 - Sailing stones, 159
 - Wheeled threats to deserts, 159
 - Readings and references, 160
- 8 Chaparral and Pinyon-Juniper Woodlands, 167**
 - Features and adaptations of chaparral, 168
 - Coastal (California) chaparral, 171
 - Chamise chaparral, 171
 - Manzanita chaparral, 171
 - Ceanothus chaparral, 171
 - Other chaparral communities, 171
 - Interior (Arizona) chaparral, 172
 - Pinyon-juniper woodlands, 172
 - Distribution and ecology, 172

- Human uses, 173
- Chaparral and fire, 173
 - Water-repellant soils, 173
 - Post-fire vegetation, 174
 - Wildlife and chaparral fires, 174
- Highlights, 174
 - Allelopathy in chaparral, 174
 - Animal associates in coastal chaparral, 175
 - Lizards and burned chaparral, 175
 - Pinyon jays, 176
- Human influences, 176
- Readings and references, 176
- 9 Montane Forests, 180**
 - Features and adaptations, 180
 - Montane Forest zones, 181
 - Lower montane zone, 181
 - Upper montane zone, 181
 - Subalpine zone, 182
 - Associated habitats, 183
 - Mountain parks and meadows, 183
 - Black Hills, 185
 - Redwoods and sequoias, 186
 - Bristlecone pine forest, 188
 - Fire in montane forests, 190
 - Highlights, 192
 - Western chipmunks and competitive exclusion, 192
 - Squirrels, bears, and pine cones, 194
 - Sky islands in Arizona, 195
 - Monarchs in winter, 196
 - Bears and moths, 196
 - Readings and references, 198
- 10 Temperate Rain Forest, 203**
 - What is old-growth forest?, 203
 - Features and adaptations, 204
 - Valleys of rain forest, 204
 - Epiphytes, canopy roots, and “scuzz”, 206
 - More about logs, 207
 - Succession on glacial till, 208
 - Highlights, 209
 - Bears, salmon, and forest enrichment, 209
 - A seabird in the forest, 212
 - Some small mammals and their ecology, 213
 - Banana slugs, 213
 - Pacific yew, 214
 - Giant salamanders and other amphibians, 214
 - Mount St Helens, 215
 - Ecological controversy, 217
 - Readings and references, 218
- 11 Coastal Environments, 223**
 - Currents and climates, 223
 - Features and adaptations, 224
 - Rocky seashores and tidal pools, 224
 - Sandy seashores, 225
 - Chesapeake Bay, 227
 - Mother Lagoon, 228
 - Submergent communities, 230
 - Seagrass meadows, 230
 - Forests in the ocean, 232
 - Oyster reefs, 233
 - Emergent communities, 235
 - Atlantic tidal marshes, 235
 - Marshes of the Gulf Coast, 236
 - Mangrove islands and thickets, 238
 - Some associated communities, 239
 - Barrier islands, 239
 - Coral reefs, 241
 - Maritime forests, 243
 - Highlights, 243
 - Synchrony at Delaware Bay, 243
 - Waterbird colonies, 245
 - A whale of a success, 247
 - Ecological challenges, 248
 - Natural disturbances, 248
 - Sea-level rise, 250
 - Readings and references, 251
- 12 A Selection of Special Environments, 259**
 - The Grand Canyon, 259
 - Caves, 262
 - Arctic ice cap, 265
 - Niagara Escarpment, 267
 - The “Father of Waters”, 268
 - The Everglades, 271
 - Fossil *Lagerstätten*: Windows into North America’s ecological past, 275
 - Burgess Shale, 275
 - La Brea tar pits, 277
 - The Florida Keys, 278
 - The Great Lakes, 280
 - Habitat highlights, 282
 - Rivers of ice, 282
 - Hot springs and geysers, 283
 - Forest in the clouds, 284
 - Granite outcrops and inselbergs, 285
 - Palm forest, 285
 - Mineral licks, 286
 - Bogs and their carnivorous plants, 287
 - Readings and references, 288
 - Appendix, 296
 - Glossary, 309
 - Index, 321

Foreword

As a professor, scientist, and student of wildlife for over four decades, I cannot think of a more important body of work to understand than ecology. This field of science is the basis for all natural history professions including anything that involves flora and fauna in its native habitat. It would be hard to impossible to understand, conserve, or manage coastal areas, deserts, forests and woodlands, grasslands, or tundra (essentially the landscapes that contribute to habitats for wildlife) and the plants and animals that occupy them without a solid understanding of ecology. Brian Chapman and Eric Bolen have brought these systems and others to life in this new edition of *Ecology of North America*.

Eric Bolen published the first edition of the book in 1998. This new edition is not just a makeover of Bolen's earlier work. Chapters were updated where needed and new information was added, including a new chapter on coastal environments to round out the landscapes examined. Another exciting addition is the inclusion of information boxes that call out special individuals and events in ecology that will be of interest to readers. Individuals highlighted include the "Fathers" of animal ecology, ecosystem science, grassland ecology, wildlife management, and prominent movers and shakers in the field such as Samuel Hearne, Rachel Carson, E. Lucy Braun, and John Muir. There are 18 information boxes scattered throughout the book with intriguing titles that are engaging, interesting to read, and informative, including: "The elk that saved a forest", "Spirit bear, the other white bear", and "Agave, margaritas, and bats". Not only are they interesting, but these selected topics are written in such a way as to make the reader to want to know more.

This book is not the typical ecology text that only gives a cursory treatment to the important landforms of North America. It is unique in that it briefly describes the basics of ecology in the first chapter, and then delves into the ecology of the major landscapes of our continent in detail. College students and layman alike can easily understand the descriptions. The writing is clear and interesting, and would be of value to anyone interested in the places they live or new places they will be visiting. For example, I have taken many visitors with me during my studies in the deserts of North America, and often hear statements like "Wow, there is nothing here – no wonder they call it a desert". It is only after visitors spend time looking at the vast amount of life and signs of life that their tunes change to one of wonder. The authors

believe that people will have a much better appreciation for the landscapes they occupy – from the fragile cryptogamic soils in deserts to freezing frogs in tundra – by reading this book prior to visiting new areas (biomes). This new volume is chock-full of interesting information including how alligators create habitat with "gator holes", introducing the reader to forests in the ocean, the importance of "scuzz" in enriching forests on land, and how *Lagerstätten* fossils and packrat middens can help humans understand their past. The writing brings together the interactions of plants, animals, and habitats (as ecology should), such as the explanations of how brown bears, salmon, and forests interact, or uses birds and thousand-year-old pines or cicadas to introduce ecological terms to readers.

The advent of computers and models has greatly changed the way students are taught and how they learn in North American universities and colleges. As more flora and fauna are reduced from living, breathing organisms to pixels on a computer screen, there seems to be less attention devoted to actual field study. Some universities even have labs in the natural sciences online. I cannot imagine learning something as basic, yet complex, as ecology from a computer, and that is clearly not the intent here. Brian Chapman and Eric Bolen bring the fascinating landscapes humans share with flora and fauna to life and also explain how mankind influences these systems.

The authors are well-known ecologists with a long history together; both have dedicated their lives to understanding ecology around the United States and passing on that knowledge to others. Because of their long-term association, the writing is seamless as though written by a single author. The book will certainly give budding ecologists and natural resource professionals a sense and better understanding of "place". The authors had a goal of following Leopold's example of emphasizing the importance of landscapes to the biologic community and to the human spirit. *Comme il faut!*

Paul R. Krausman
Certified Wildlife Biologist®
Fellow and Past-president of The Wildlife Society
Emeritus Professor of Wildlife Ecology,
University of Arizona
Boone and Crockett Professor of Wildlife Conservation,
University of Montana

Preface

About this book

Ecology of North America stems from our belief that many college students – and non-students – lack much awareness about the natural world in which they live. Thanks to the effects of mass media, some may indeed deplore the destruction of far-off tropical rain forests or the plight of whales; however, too few are acquainted with the all-but-vanished inland sea of tallgrass prairie, the ‘gator holes of the Everglades, or the enigmatic population cycles of lynx (and other) animals in the spruce-fir forests of North America here at home. Our conviction rests, collectively, on more than 60 years of college teaching and countless public presentations. Moreover, until the first edition appeared no single text designed for classroom use had attempted to survey the ecological diversity of North America’s vast landscape. The first edition of *Ecology of North America*, drafted while one of us (EGB) taught a course of the same name, was therefore conceived and eventually born as a tool fashioned for undergraduate instruction.

Shortly after the first edition appeared preparation began for a second edition, but various circumstances precluded its completion (not least EGB’s retirement). Hence, if a new edition was to materialize, a coauthor was needed to revitalize the work. BRC then entered the scene when, after careers at other universities, he joined the faculty at UNCW as provost and professor of biology. Notably, his teaching and research experiences paralleled the basic themes expressed in *Ecology of North America*, and a marriage of interests thereafter followed. Our association in fact began decades ago when BRC was completing his doctoral work at Texas Tech University, where EGB was then serving on the faculty and as a member of his advisory committee. This edition is therefore the result of a long-standing partnership founded on both a deeply shared interest in plant and animal ecology and a collegial and personal friendship.

As in the previous edition, the geographical coverage begins in the far north (the frigid tundra), continues southward into boreal and deciduous forests, swings westward into the interior grasslands and deserts, and continues to the western mountains before traveling onward to chaparral and temperate rain forests. Sites along North America’s long and varied coastline conclude our regional tour. We present a selection of unique areas (e.g., the Grand Canyon), expanded in this edition to include the fossil history offered by the La Brea tar pits and Burgess Shale. We also visit far-flung locations such as the Polar

Ice Cap, Mississippi River, and Florida Keys, as well as stopping to investigate pitcher plant bogs and other habitats of interest on the way. As appropriate to a new edition, we have supplemented and updated the previous text with a variety of subjects ranging from cicada cycles and the moth-eating habits of bears, to the mysterious disease now devastating bat populations and the havoc wrought to the Everglades by introduced pythons.

Some entirely new features highlight the second edition including a chapter dedicated to coastal environments, among them seagrass communities, tide pools, and barrier islands, and ecological portraits of Chesapeake Bay and the Laguna Madre. The new chapter also describes phenomena such as the vital relationship between horseshoe crabs and migrant shorebirds. “Infoboxes” likewise appear for the first time; these present stand-alone descriptions, including biographies, as disparate as spirit bears, Lucy Braun, and the conservation successes of gray whales and Maine’s puffin colonies. To our delight, color photos now complement the gallery of black and white images.

Each chapter ends with “Readings and references,” some of which present basic descriptions of community structure. These appeared early in the development of ecology, yet still provide an essential foundation for many readers. Others sources which have been published more recently represent new discoveries and refinements of previous concepts, such as: revelations about the hunting success of spirit bears; evidence of inbreeding depression in Isle Royale’s isolated wolf population; the biotic community in the canopy of towering redwoods; and expanding knowledge of the beneficial impacts of nutrients derived from salmon carcasses. We also list works of greater scope for those seeking deeper insight into the subjects at hand.

Our focus

This edition, updated with both new and expanded coverage, broadly targets an audience of both undergraduate students and the general public. In doing so, we continue writing in a “user-friendly” format that appends Latin taxa and literature sources instead of embedding these within the text in the style of scientific journals. Understandably, some professional biologists may fault our format, but we aim to hold the attention of readers who may place higher priorities on the primary message of the book. For the same reason, we shied from marching lockstep through a catalog of species, soil types, and

weather regimes for every unit; these are provided only when they seem important. Otherwise, our steady focus is on key or at least interesting plants and animals and their interactions.

Finally, we were guided by the thoughts of Aldo Leopold who, in *Sand County Almanac* (1949), entwined three thoughts into a common theme: “land is a community” that forms the fabric of ecology; land should

be cherished and respected as an “extension of ethics”; and “land yields a cultural harvest.” Leopold wedded science, ethics, and aesthetics. In doing so, he established an understanding of nature that we, in our own small way, have tried to nourish.

Brian R. Chapman, Huntsville, Texas
Eric G. Bolen, Wilmington, North Carolina

Acknowledgments

Our task would surely have foundered without the generous advice, knowledge, and contributions of many colleagues. Merely to list their names seems far too superficial, but we are nonetheless indebted to all, including anyone we have inadvertently omitted.

Our continued thanks are extended to those who contributed to the first edition (and in some cases, this edition as well): C. Davidson Ankney, Michael J. Armbruster, Stephen F. Arno, Guy A. Baldassarre, Thomas C. Barr, Jr, Mark Blumenthal, Carlton M. Britton, Lincoln P. Brower, C. Alexander Brownlow III, Dana C. Bryan, George C. Carroll, Donald D. Clark, Robert T. Coupland, John A. Crawford, James E. Deacon, Leonard F. DeBano, Ronald I. Dorn, Harold E. Dregne, Christine S. Dutton, Ellen Faurot-Daniels, Maeton C. Freel, Peter M. Frenzen, Ron R. George, Paul L. Gersper, Paul N. Gray, Lance H. Gunderson, Craig A. Harper, Wendy Hodgson, John R. Holsinger, Jon E. Keely, Fritz L. Knopf, Paul R. Krausman, Melissa Kreighbaum, Douglas W. Larson, Gordon S. Lind, Erik Lindquist, Glenn Longley, Charles Lowe, Richard Mack, John Mangimeli, Paul S. Martin, Roger Martin, Joseph R. McAuliffe, Burruss McDaniel, Robert W. Mitchell, Robert. H. Mohlenbrock, Elizabeth M. Morgan, Fritz P. Mueller, Richard Murzin, Nalini M. Nadkarni, S. Kim Nelson, David J. Odz, Karl A. Perry, James Petterson, Linda Pin, Donald Pinkava, Thomas L. Poulson, Donald R. Progulske, William O. Pruitt, Jr, John Riley, William L. Robinson, William H. Romme, Donald H. Rusch, Susan Shultz, John G. Sidle, John C.F. Tedrow, Jack W. Thomas, Walter M. Tovell, Merlin D. Tuttle, Gary Valentine, Thomas R. Van Devender, B.J. Verts, Doug Waid, Frederic H. Wagner, John R. Watson, Ronald R. Weeden, Nathaniel R. Whitney, Edward O. Wilson, M. Eugene Wright, Paul J. Young, and Paul Zedler.

For their contributions to the second edition, we also thank Kathy L. Allen, Donna Anstey, Angie Babbitt, Candice Bressler, Sanford Brown, Lisa D. Chapman, William R. Clark, Sherry Cospser, Tara Cuvelier, Pamela Dibble, John Dokken, Catherine Evans, Debbie R. Folklets, W. Mark Ford, Eric Hallerman, Richard W. Halsey, Thomas Henry, Susan Jacobson, Rick Kesterman, Jack L. Kindinger, Roel Lopez, Jim Lovett, D. A. Mascarelli, Joyce Maschinski, Vincente Mata, Laurie McBurnie, Danny L. McDonald, Thomas C. Michot, Thomas E. Moore, William F. Moore, Ashley R. Morgan, Paige A. Najvar, T. J. Pernas, Luis Prado, Kristen Quarles, T.P. Quinn, W. Scott Richardson, Celia Rozen, Charles H. Smith, Mike Smith, Autumn J. Smith-Herron, Clyde Sorenson, Nancy P. Stewart, James Stubbendieck, Orly

(Chip) Taylor, Monte L. Thies, Nancy Thompson, Jace Tunnell, David Welch, Josh Westerhold, Don White, Jr, Richard Whyte, and Kathy S. Williams.

Colleagues at the University of North Carolina at Wilmington generously provided advice and assistance as one or both of the editions of the book were being prepared. Special mention goes to W. Franklin Ainsley, Walter J. Biggs, Thomas F. Britt, Lawrence B. Cahoon, Troy M. Clites, Mike Durako, D. Wilson Freshwater, Jocelyn Gaudet, Gilbert S. Grant, Melissa Grey, Neil F. Hadley, William B. Harris, Paul E. Hosier, John R. Huntsman, Stephen T. Kinsey, David L. LaVere, David G. Lindquist, John J. Manock, R.D. McCall, Ian A. McLaren, William McLellan, Daniel W. Noland, D. Ann Pabst, David E. Padgett, Beth A. Pardini, James F. Parnell, Joseph R. Pawlik, Martin H. Posey, David J. Sieren, Amanda Southwood, Peter N. Thomas, Carmelo Tomas, W. David Webster, and Barbara L. Wilson. The staff at the Randall Library at UNCW expertly – and patiently – accommodated requests for often obscure bibliographic information and materials.

Associates at Sam Houston State University, the University of Houston-Downtown, and colleagues at several other universities thoughtfully provided essential information, assistance with photographic needs, and various kinds of advice and support during preparation of the second edition. Among these, special thanks go to Loren K. Ammerman, Jerry L. Cook, Quenton R. Dokken, Richard Eglsaer, Jaimie L. Herbert, David R. Hoffpauir, William I. Lutterschmidt, Lisa A. Shipley, Melissa S. Sisson, and John W. Tunnell, Jr. Sandra S. Chapman painstakingly proofread the manuscript, and we appreciate her keen eye and invaluable suggestions.

Persons and/or agencies providing photographs are acknowledged in the appropriate figure captions, as are the artists who prepared maps and line drawings; special thanks go to Tamara R. Sayre. Many of her illustrations appear again in the second edition as testimony to her talent and, certainly, to the importance of her work to the content of the book.

We benefitted from constructive comments provided by colleagues who reviewed the first edition as well as those who examined draft chapters prepared for the second edition, among them Leonard A. Brennan, Paul R. Krausmann, Bruce D. Leopold, Joyce Maschinski, Eric Ribbins, Fred E. Smeins, and an anonymous reviewer.

We are especially indebted to Paul R. Krausman who provided advice and constructive suggestions during the

preparation of both editions, and who graciously wrote the Foreword for the second edition.

Rachael Ballard, Executive Commissioning Editor at John Wiley and Sons, unhesitatingly authorized this project, and we enjoyed working with her and Delia Sandford, Managing Editor. We also thank Fiona Seymour, Senior Project Editor, and Audrie Tan, Project Editor, who guided the manuscript through the start-to-finish maze of production. We owe a special debt of gratitude to our copyeditor, Elaine Rowan, whose thoughtful

improvements made what we were attempting to say far more intelligible.

Clearly, many have contributed, but we alone remain responsible for any errors that escaped unnoticed.

Finally, our wives, Sandy and Elizabeth, endured more neglect than we care to admit while this project inched toward completion. We marvel at our good fortune to find such wonderful, loving, and supportive partners. To them, we dedicate this book (albeit that it seems so little in return for so much).

CHAPTER 1

Introduction

In all things of nature there is something of the marvelous.

Aristotle

Think big for a moment. Imagine a transect running north to south, stretching across the midpoint of pristine North America in the year 1500. We will use the 100th meridian, which traces the right-hand edge of the Texas Panhandle, as our line of travel. At its northern end – the North Pole – the transect begins in a desolate cap of ice and snow and then crosses a vast Arctic landscape of tundra before traversing a wide band of spruce and fir known as the Boreal Forest. Wolves howl in the distance; the scattered remains of their most recent kill – a sickly moose – lie still fresh atop the deep snow. South of these dark forests spread the plains, grasslands grazed by millions of bison and even larger numbers of prairie dogs. After leaving the plains, our excursion takes us across the narrow, clear rivers and low hills of the Edwards Plateau before dropping into a region studded with low, thorny trees. The transect ends, for our purposes, when it reaches northern Mexico and the yucca-spiked Chihuahuan Desert.

A similar transect running east to west across the waist of North America at first encounters the sandy beaches and dunes of the Atlantic seashore. Here, at the latitude of Virginia, the transect crosses what once was an immense forest of oak and hickory (Fig. 1.1). According to folklore, an ambitious squirrel could have journeyed from the Atlantic seaboard to the Mississippi River 300 years ago without the necessity of ground travel, false testimony to the idyllic notion that an unbroken canopy of deciduous forest once stretched across more than a third of North America.

Westward, with the Appalachians, Cumberland Plateau, and Mississippi River behind us, the forest gradually thins and our transect enters the grasslands. Our trek skirts the southern edge of the Prairie Pothole Region – North America’s famed “duck factories” – and pushes onward toward the plains. As we cross our north-south transect in western Kansas, courting prairie chickens dance and boom in the background and

black-tailed jackrabbits crouch, ears lowered, to escape our notice. Still farther west rise the Rocky Mountains with their rather distinctive zones of vegetation, after which we enter into a desolate terrain of sagebrush in the Great Basin. In the distance loom the peaks of the Sierra Nevada where the largest of trees, the giant sequoia, almost defy description. In a quiet grove of these immense trees, we might allow our imagination to behold the ghost of John Muir (1838–1914), the hard-trekking Scot who championed wilderness preservation.

By the time we reach the rocky seashore of the Pacific Ocean, our transect will have dropped into the Central Valley of California and then risen upward to cross the Coastal Range that rims the western edge of the continent. Offshore, frisky sea otters float above submerged kelp forests. North of where we stand are the old-growth forests of Sitka spruce and western hemlock, heavily draped with epiphytes, replete with spotted owls and maples. On the Alaskan coast, vulnerable hoards of migrating salmon attract giant bears to the rushing streams each year. To the south are chaparral-covered hills, and beyond these are the beckoning Joshua trees in the Mojave Desert. Still farther south at a small site in the mountains of Mexico are firs cloaked each winter with millions of slumbering butterflies. Our telescoped journey across North America has been brief to be sure, but perhaps it is long enough to preview the contents of this book.

A brief overview of ecology

Ecology is the branch of biology that investigates the inter-relationships between organisms and their environment. The original name *oekologie*, based on the Greek word *oikos* meaning “home”, was coined in 1866 by German zoologist Ernst Haeckel (1834–1919). An ecological study of any species involves a detailed examination of an organism’s life history and biological requirements, the physical



Figure 1.1 A vast expanse of eastern deciduous forest once covered much of North America east of the Mississippi River, but the romantic notion of an unbroken canopy 300 years ago is inaccurate. Indeed, Native Americans and natural processes, such as fire, periodically cleared many areas, some quite large, within the primordial forest. A mountaintop bald, visible atop the Smoky Mountain ridge (right), represents a natural opening described in more detail in Chapter 3. Photo courtesy of Brian R. Chapman.

environment in which the organism lives, and its interactions with the other species that occupy the same area. Physical features of the environment (e.g., temperature, soil type, and moisture) influence the distribution and abundance of organisms, but all living things process materials from the environment and transform energy as they grow and reproduce.

The ecosystem

In 1935, English botanist Arthur G. Tansley (1871–1955) proposed the term **ecosystem** to characterize the flow of energy and matter through a network of **food chains** collectively known as a **food web**. Energy passes through a series of **trophic levels** (i.e., feeding levels), the functional parts of an ecosystem. These rest on a foundation of non-living matter, the **abiotic level**, which consists of air, soil, and water. When fueled by sunlight, the abiotic level provides the fundamental components required for **photosynthesis** by species known as **primary producers**, which are represented by green plants such as algae, grasses, and trees and first in the series of trophic levels. **Primary consumers** – rabbits, deer, or other **herbivores** that consume the energy and matter bound in green plants – represent the next trophic level, followed by **secondary consumers**; the latter are predators such as foxes or hawks. **Tertiary consumers**, sometimes known as apex predators, are represented by animals such as polar bears or mountain lions. The final trophic level, **decomposers**, is populated by scavengers, bacteria, and fungi that return the tissues of dead plants and animals to their elemental form (Fig. 1.2). An ecosystem is therefore an area or volume in which energy and matter are exchanged between its living and non-living parts.



Figure 1.2 In forests decomposition is commonly thought to originate with logs, but it often begins when a dead tree remains upright as a snag. The bracket fungi on this snag possess enzymes that break down lignin and other complex chemicals found in wood, and the fungi absorb the nutrients for their growth and reproduction. When the snag eventually falls, other decomposers on the forest floor will return its elements to the soil. Photo courtesy of Brian R. Chapman.

Abiotic limits

Many environmental influences – such as moisture, temperature, the availability of nutrients, wind, and fire – limit the kinds and abundances of organisms that populate an ecosystem. In 1840, German organic chemist Justus von Liebig (1803–1873) was the first to recognize the role of abiotic limitations in nature. After studying the relationships between surface soils and agricultural plants, Liebig concluded: “The crops of a field diminish or increase in exact proportion to the diminution or increase of the mineral substances conveyed to it in nature.” Liebig recognized that each plant requires not only certain minerals, but each must also be present in the proper quantity for the plant to flourish. If a required

nutrient is absent, the plant will not survive. Moreover, if the essential food substance is present only in a minimal amount, the plant’s growth will be correspondingly minimal. In later years, this concept became known as the **law of the minimum**.

Later studies indicated that the growth and survival of living organisms also may be limited by an overabundance of a required substance (e.g., terrestrial plants require moisture, but die when waterlogged for a prolonged period). Plants and animals are successful only when they live in an environmental range between too much and too little, that is, within their limits of tolerance. Victor E. Shelford (1877–1968; Infobox 1.1) incorporated the concept of maximum

Infobox 1.1 Victor E. Shelford (1877–1968), Father of animal ecology

The scientific career of Victor Shelford began in 1899 when he enrolled at West Virginia University where his uncle, the assistant state entomologist, influenced his lifelong interest in insects. In 1901, however, the president of the university moved to the University of Chicago where he offered a scholarship to Shelford who accepted and eventually earned a Ph.D. (in 1907) dealing with tiger beetles and dune vegetation on the shores of Lake Michigan. Likely for the first time, this study associated animals with the successional changes in a plant community, a concept pioneered by his academic mentor Henry C. Cowles (1869–1939). Shelford thereafter joined the faculty at Chicago but moved to the University of Illinois in 1914, where he spent the remainder of his academic career.

In 1913, Shelford published his first book, *Animal Communities in Temperate America*, a landmark of its day. He helped found the Ecological Society of America and, in 1916, became its first president.

Along with his university duties, Shelford served as the laboratory supervisor for the Illinois Natural History Survey during 1914–1929 and, during alternate summers between 1914 and 1930, as director of marine ecology at the Puget Sound Biological Station. His research interests included topics ranging from benthic communities in both freshwater and marine environments to lemming populations in Arctic tundra. His experiments covered the physiological and behavioral responses of animals to temperature and other variables in climate-controlled chambers. In the field, he used photoelectric cells to determine light penetration into seawater. His research, which often employed novel equipment, led to a manual entitled *Laboratory and Field Ecology* (1929). On the practical side, Shelford also studied termites and other insect pests, as well as the response of fishes to sewage treatment. He eventually reduced his lab work in favor of spending more time on field studies, especially those that concerned food chains, structure, and other analyses of natural communities.

In the 1930s, Shelford began collaborating with Frederic E. Clements (1874–1945), a prominent plant ecologist. Their relationship was cordial but strained at times as, much to Shelford’s dismay, Clements harbored reservations about the role animals played in vegetational development. Nonetheless, in 1939 the two produced *Bio-Ecology*, a book that integrated plants and animals into the formation of communities.

Shelford’s marathon field trips were legendary experiences for students; some included several weeks of camping while visiting prairie, forest, desert, or tundra. Indeed, he wanted his students to study first hand every major biome in North America. Reelfoot Lake in Tennessee, created by a cataclysm of earthquakes in the winter of 1811–1812, was a regular stop for many years; the site included opportunities to study oxbow lakes, cypress sloughs, and floodplain forests. Wherever they went, however, Shelford steadfastly donned shirt and tie in the field. Shelford’s career represented a major link in a chain of renowned ecologists that began with Cowles and continued with Shelford, followed by his student S. Charles Kendeigh (1904–1986) and, in turn, the latter’s student Eugene P. Odum (1913–2002), who is regarded as the founder of modern ecology.

Shelford retired from university life in 1946, but he hardly remained inactive. His passion for preserving natural areas with fully intact communities initiated the Ecologist’s Union, which evolved into *The Nature Conservancy* whose mission is now global in scope. His association with Clements helped Shelford develop his appreciation for the concept of biomes and, after years of work, resulted in a monumental treatise, *Ecology of North America* (1963), the inspiration and namesake for the book at hand.

Soon after Shelford completed his graduate studies at Chicago, he was advised by a prominent scientist to “discontinue this field of ecology” and instead pursue biology with a traditional focus on individual organisms. Fortunately, Victor E. Shelford chose otherwise.

and minimum limits on environmental condition into the **law of tolerance** in 1913. Some organisms are capable of living within wide ranges of conditions of one or more environmental factors, whereas others have narrow limits of tolerance. Certain species are capable of acclimatizing to different environmental limits as seasons or conditions change.

To complicate things further, an organism may tolerate a wide range of conditions for one environmental factor, but be restricted by a narrow range for another. In the latter case, the distributional range of the species will be restricted by the condition for which it possesses the narrowest tolerance.

Climate and topography

The climatic conditions of any region are determined by many factors, including latitude, seasonal temperature ranges, the amount and periodicity of rainfall, and location on the continent (e.g., interior versus coastal). Geographical features, such as proximity to mountain ranges or bodies of water, can alter local conditions enough to influence the type of plant associations that can exist. Thus, vegetation maps of North America (Fig. 1.3) illustrate the influence of climate. The northern region of North America has two broad, east–west belts of natural vegetation: tundra and boreal coniferous forest. These follow a gradient correlated with temperature patterns, but the vegetative zones on the remainder of the continent are more closely associated with the availability of moisture. Precipitation decreases from east to west while evaporation, which is largely influenced by temperature, increases; this interaction results in a series of north and south vegetation bands. Along the coasts, regions of higher humidity support forests. The broadest zone of forest is on the eastern half of the continent where rainfall exceeds evaporation.

Mountains intercept winds and directly influence regional climates. As winds ascend a mountain slope, the air mass cools and gradually becomes saturated; cool air holds less moisture than warm air. The windward sides of mountains usually receive rain at mid- to higher elevations as the moisture is, in effect, wrung out of the air. On the leeward side, descending cool dry air absorbs moisture from the soil and vegetation creating **rain shadow**, an area where evaporation greatly exceeds precipitation. The Great Basin Desert on the eastern (leeward) side of the Sierra Nevada Range illustrates the impact of a rain shadow, whereas the western slopes of the mountains support lush forests.

Topography and vegetation create local environmental conditions that can differ substantially from the overall climate of a region. For example, the **microclimate** under a dense clump of bushes does not share the same temperature, moisture, and wind conditions as may be found in an open area just a short distance away. Similarly, the north-facing slope on a mountain does not

receive the same solar radiation as a south-facing slope. Because of the difference in exposure, the evaporation rate on north-facing slopes may be 50% lower, the temperature is lower, and soil moisture is higher. Thus, two sides of a mountain, even at the same elevation, are often occupied by different compositions of plant and animal species.

Soils and soil profiles

Climatic features, especially temperature and precipitation, influence the formation of soils. Soil is a complex mixture of minerals, organic matter, water, and air, forming a substrate harboring bacteria, fungi, and other small organisms. Minerals are derived from the weathering of parent material, usually rock, by chemical and physical processes. The microorganisms (e.g., decaying organisms) contribute to soil formation and development by breaking down organic matter, fixing atmospheric nitrogen, and contributing to nutrient cycling.

The US Department of Agriculture (USDA) recognizes twelve textural classes of soil based on various compositions of three particle sizes of minerals – clay, silt, and sand – but soil also may contain larger mineral particles ranging in size from pebbles to large fragments of rock. Of the textural classes, loam (composed of varying percentages of clay, silt, and sand) is the most valuable because it supports many agricultural crops. The USDA also developed a soil classification system that places all the soil types in the world into twelve **soil orders**. These soil groups are defined using a set of characteristics that includes texture as well as physical and chemical properties. The name of each soil order ends in *-sol*, which is derived from the Latin word *solum* meaning “soil”. For example, about 16% of the world’s surface is covered by entisols which have moderate to low fertility; mollisols, which are the most productive agricultural soils, only occur on about 7% of the Earth’s surface.

Anyone who has dug a posthole or planted a tree knows that soils are organized into distinct layers or **horizons**, descending from the ground surface down to the bedrock. These layers form a **soil profile**, and are the product of weathering and the actions of vegetation, temperature, rainfall, and microorganisms acting for millennia on parent material in a specific locality. The organic material – leaf litter and decaying plant and animal matter – on the ground surface forms the **O horizon**. The O horizon is generally thin or absent in deserts and grasslands, but may be quite deep in forests. The first layer of soil, the **A horizon**, is often called “topsoil” and is usually rich in **humus**, the dark-colored products of decomposed organic materials. The A horizon is absent in most deserts, but can be up to 1 m (3 feet) deep in the fertile wheat-growing prairies of Washington State. Progressively downward in the soil profile, the E, B, and C horizons represent zones where the vertical processes of leaching gradually reduce the




Figure 1.3 The predominately east–west distribution of major biomes in North America is determined largely by precipitation. In contrast, the north–south arrangement of major plant associations in the middle and southern regions of the continent result more from interactions of temperature, evaporation, and elevation. Illustration courtesy of Tamara R. Sayre and Brian R. Chapman, based on Brown et al. (2007).

organic content of each successive layer, thereby increasing the amount of mineral matter. Bedrock is reached at the R horizon, which typically represents parent material.

Biotic community

A **biotic community**, the living part of an ecosystem, is composed of many species, but the number of species and their relative abundance diminish “upward” through a food chain. More species of plants exist in the producer level than do herbivores functioning as primary consumers. Likewise, herbivores are more numerous than the carnivores in the upper trophic levels. These relationships result in a pyramid-shaped community, with numerous (both in species and abundance) green plants at its base, fewer herbivores, and fewer still **carnivores** at its apex (e.g., mountain lions). In Chapter 3, we will describe a relatively simple pyramid of vegetation, moose, and wolves on Isle Royale.

Community succession

Plant **succession** is the relatively predictable sequence of vegetational development within a geographic region. The composition of the local fauna also changes as the vegetation develops and, to express the full ecological impact, the process is best regarded as biotic succession (i.e., sequential changes in both plants and animals). In practice, ecologists usually refer to the concept simply as succession and use the term more in reference to plants than to animals.

Notions about succession are embedded in several 19th century sources, including discourses of Henry David Thoreau (1817–1862) who described the regrowth of forests near Concord, Massachusetts. The first ecologist (at least in North America) to describe succession was Henry C. Cowles (1869–1939), who based his observations on the changes in dune vegetation on the shores of Lake Michigan. Later, Frederic Clements (1874–1945) proposed a complex hierarchy to describe the development of vegetation over time, but only a few of his terms remain widely used today.

Succession is currently viewed as a process of multiple invasions. As certain species replace others because of their own particular **adaptations** (e.g., tolerance to shade in an early growth stage, or greater ability to disperse), the composition of the community changes. The process of succession is also strongly influenced by local environmental factors such as soil type and the availability of water or certain nutrients.

Plant succession: from pioneer to climax

To perceive succession visually, consider a cleared landscape such as an agricultural field in a region with a moderate climate. Left unplowed, the exposed soil is

soon covered with so-called weeds – dandelions and asters are typical invaders in many areas – that represent **pioneer** vegetation. Such herbaceous immigrants have three general characteristics: (a) they thrive on disturbed soils where they tolerate harsh conditions (e.g., high soil temperatures, limited moisture and, in many cases, few nutrients); (b) they produce large numbers of seeds with adaptations for widespread dispersal and rapid germination but, when necessary, their seeds can persist in the soil for long periods; and (c) they are commonly, but not always, annuals. Some ecologists regard these as “opportunistic species”, plants that quickly take advantage of opportunities to colonize (or re-colonize) sites where competitive species are absent. The pioneer community accordingly consists of relatively few species; recently abandoned fields, for example, are often covered by blankets of single species.

Eventually, the pioneers give way to perennial species, which include plants with established roots systems, storage organs, and defensive structures. These species are better competitors and they eliminate the pioneers. Broomsedge and goldenrod are representative species. Shrubs soon appear, to be replaced later by trees. In many locations, a pine forest develops first, followed by a forest of oaks or other deciduous hardwoods. Of course, the species in these sequences vary by region, soil type, and other factors, but the pattern is consistent.

The final stage in succession is known as the **climax**, a term ecologists use to characterize regional vegetation. Climax communities, such as the Oak–Hickory Association in the Eastern Deciduous Forest, are mature ecosystems consisting of characteristic plants as well as animals (e.g., wild turkeys and eastern gray squirrels). When compared to pioneer and other successional stages, climax communities are (a) neither as **hydric** (wet) or **xeric** (dry) as the earlier stages and instead represent **mesic** conditions; (b) more complex and better organized; they have complicated food webs and more interspecific relationships; (c) include more species, which tend to be relatively large and long-lived and have low reproductive rates; and (d) comparatively stable (e.g., resistant to invading species).

Primary and secondary succession

Old-field succession on abandoned farmlands is a real-world event often described to illustrate the sequence of community replacements. Fallow fields, which were widely available in the late 1930s, are useful sites to study succession because fields in various stages of regrowth are available (i.e., the date of abandonment, and hence the “age” of each field, and its vegetation can be determined from court records). One of the earliest studies of old-field succession on an abandoned farm in the Piedmont region of North Carolina revealed a sequence of herbaceous pioneers such as aster and ragweed, followed by the arrival of pine between years 5

and 15, and then full development of an oak–hickory forest 150 years after the fields were last cultivated.

Old-field succession illustrates **secondary succession**, the recovery of a previously vegetated but disturbed site. Secondary succession also occurs after fires remove the previous vegetation. These sites generally have well-developed soils that can support a variety of plants. The abandoned fields in North Carolina described in the previous paragraph were later cleared of forest vegetation and cultivated for many years thereafter. **Primary succession**, in contrast, occurs at sites devoid of previous vegetation, examples of which include dunes, atolls, volcanic cones, and land exposed by retreating glaciers. Such sites typically lack mature soils and have few nutrients. Primary succession, because it usually requires soil building, spans long time periods before the climax stage is reached.

Climax vegetation, once established, should not be viewed as static. Instead, it experiences a state of dynamic equilibrium, that is, it continually replaces itself. A climax community will maintain its composition so long as the climate remains stable and there are no major disturbances such as fire that return the community to an earlier stage of development. In some places, fire recurs often enough to inhibit the development of a climax community; such a setting is said to be a **subclimax** community.

Succession and species abundance

The number of species in a community increases steadily as succession progresses. Because of their harsh environment, pioneer communities typically contain only a few species but in large numbers. A barren field, for example, receives the full impact of solar radiation during the day creating surface temperatures ill-suited to, or even fatal for, many organisms, but at night the barren field cools rapidly due to a lack of both solar radiation and an insulating blanket of vegetation. The bare soil is also exposed to the full force of wind and rain, often creating unstable conditions (e.g., erosion). Consequently, only a limited number of species are suited to such environments and those few form the pioneer community. Indeed, pioneer communities are sometimes monotypic in their composition.

Physical conditions become less harsh after pioneer vegetation gains a foothold. The herbaceous blanket ameliorates extremes in temperature, reduces runoff, and discourages wind erosion, enabling the soil to hold more moisture. Other plants find suitable conditions and the number of species, including insects, rodents, and other animals, steadily increases during the middle stages of succession.

While it is tempting to conclude that climax communities contain a maximum number of species, the peak is reached slightly before the climax reaches its full development. This relationship results from heightened biological competition, that is, some less-competitive

species (e.g., goldenrod) are eliminated when the dominant climax species (e.g., oaks) reach their fullest development. The relative degree of species reduction at climax is somewhat greater in forests and other mesic communities than in communities where the physical conditions are limiting (e.g., Arctic Tundra).

The biome concept

Climax vegetation with similar characteristics develops in broad areas throughout the world. Although the species may vary, the vegetation over a large region exhibits a common appearance and structure (i.e., **physiognomy**) in response to analogous physical environments. Because of their similarities, climax vegetation throughout the world may be grouped into one of several ecological units, each known as a **biome**. Thus, the grasslands in North America (Great Plains), Eurasia (steppes), South America (pampas), and Africa (veld) are united in the Grassland Biome. Similarly, the northern evergreen forests that form a band across North America and Eurasia form the Boreal Forest Biome. Of the ten terrestrial biomes that are generally recognized worldwide, the only other biomes represented in North America include Desert, Temperate Deciduous Forest, Temperate Rain Forest, and Tundra. Despite the primary focus on their respective climax communities, however, each biome is best viewed as a mosaic that includes areas of early and mid-successional communities resulting from various types of disturbances. For example, stands of longleaf pine, although clearly not representative of broad-leaved deciduous trees, represent a significant subclimax community within the Eastern Deciduous Forest.

Animals should of course not be overlooked when considering biomes but, by tradition, biomes are named to reflect their dominant vegetation. In the past, “Spruce–Moose” was proposed to identify the Boreal Forest Biome, thereby including a major animal in the name, but this and similar designations never gained acceptance. Moreover, biomes are named for climax species, which is not the case for moose (as noted in Chapter 3).

Biodiversity

Biodiversity is shorthand for biological diversity and refers to the number of species that naturally occurs in a defined space or ecological unit, taking into account the relative population size of each species. **Species richness** is a similar term, although it does not account for the relative abundance of each species. The ever-growing human population and accompanying consumption of resources place increasing pressure on the world’s fauna and flora. As a result, biodiversity is declining. Cities and other developed areas are obvious examples of locations where biodiversity has become severely limited. The destruction of tropical rain forests eliminates large

numbers of species, but many other human activities (e.g., acid rain) also reduce biodiversity in places where concrete, bulldozers, or chainsaws are not in evidence.

The “species richness gradient”

The correlation between number of species and latitude is one of the more intriguing features of biogeography. Alfred Russell Wallace (1823–1913), who discovered **natural selection** independently of Charles Darwin (1809–1882), was among the first to highlight that “animal life is, on the whole, far more abundant and varied within the tropics than in any other part of the globe.” This pattern, which applies equally to both plants and animals, begins with the incredible number of species in the tropical rain forests at the equator. North or south of the equator, species richness gradually declines toward the poles, where biotic communities consist of a rather small number of species. In the Northern Hemisphere, a steady reduction in the number of nesting birds is reflected at intervals along a gradient northward from the tropics: Columbia (1400 species); Panama (1100); Guatemala (470); New York (195); Newfoundland (81); and Greenland (56 species). A gradient of similar magnitude occurs in the biota of the Southern Hemisphere, reaching an extreme in Antarctica where terrestrial vertebrates and vascular plants are completely absent.

What might explain this pattern? Some short answers follow, but the debate continues. First, tropical climates are relatively stable year round and offer favorable conditions for life (i.e., warm and humid), whereas seasonal differences become steadily more apparent at higher latitudes. Tropical regions are also relatively free of dramatic disturbances. Conversely, disturbances such as drought and hard winters and, over the long term, glaciations occur with greater frequency in the middle and higher latitudes. Such disturbances hinder the course of diversification. While the influence of abiotic factors is lessened in a stable climate, biotic interactions – especially competition, herbivory, and predation – increase in importance. Under these conditions, tropical regions become a “diversity pump” where resources are partitioned, **speciation** increases, and **extinction** decreases. For example, as predation increases, the number of individuals in a prey population decreases, which enables a greater number of other species to coexist. At higher latitudes, however, the highly variable physical environment limits specialization and results in fewer species.

Time is the second factor to interact with climatic stability. Because diversity is the product of **evolution**, greater diversity results when and where evolution can proceed uninterrupted for long periods of time. Long periods of stability provide areas such as tropical rain forests and coral reefs with adequate time to produce rich biotas. Elsewhere, however, the regions today occupied by Tundra and Boreal Forest biomes only recently (from

a geological perspective) emerged from a glacial blanket and have had much less time to evolve a mature biota.

Abundance and availability of food also contributes to species richness along a latitudinal gradient. In tropical regions, for example, plants of one kind or another bear fruit all year round, thereby providing an abundant and consistently available food source for **frugivorous** species. These in turn become the food base for various predators and scavengers. Conversely, fruiting is markedly seasonal in temperate regions, and there are fewer frugivores. Such a relationship is evident when the species diversity of bat fauna is compared by latitude. In temperate areas, bats feed on insects and must either migrate or hibernate when winter limits their food supply; in tropical areas, where both fruit and insects are abundant and available all year round, bats can specialize and more species have evolved. A correlated factor is the increase in primary productivity along a gradient from pole to equator; as solar radiation increases, so does plant production. Consequently, more consumers can exist which encourages competition and other biological interactions (see above). Interestingly, the latitudinal pattern for terrestrial species is reversed for marine mammals, but nonetheless remains based on food abundance. Species diversity of baleen whales and carnivorous pinnipeds (e.g., seals) is greater at higher latitudes where the abundance of their prey is directly or indirectly related to an immense base of planktonic foods.

The pattern in which food or other resources is distributed within a landscape, referred to as patchiness, also seems to be involved. Greater patchiness, typical in the tropics but much less so in the Boreal Forest or Arctic Tundra, may increase diversity by accelerating the development of subspecies and species among isolated populations. Habitat patches promote isolation, and isolation promotes speciation.

To summarize, terrestrial species diversity shows a global pattern of increase from the poles to the equator. In terms of the North American biota, a band stretching from Labrador to Alaska contains far fewer species than a band of the same width stretching from Virginia to California. Various factors contribute to this relationship including environmental stability, biological interactions, solar energy, and patchiness.

Biodiversity “hotspots”

Biodiversity is widely regarded as a global resource, yet extinction rates soar as the march of expanding human populations steadily degrades natural habitats. Some conservationists estimate a species is lost every day while others believe the rate is nearly one per hour. To help stem this destruction, some 25 areas around the world with important but endangered environments – notably those with large numbers of **endemic** species – have been designated as biodiversity hotspots. Each hotspot



Figure 1.4 Madrean pine–oak woodlands, which occur in deep canyons and remote sky-islands in the rugged mountains of northern Mexico and the southwestern United States, represent sites with an amazing diversity of flora and fauna. Usually surrounded by arid habitats, these isolated hotspots – this one in Hell’s Canyon in the Davis Mountains, Texas – illustrate an insular type of distribution. Photo courtesy of Brian R. Chapman.

(a) contains vegetation that includes at least 0.5% (or alternately 1500) species of endemic vascular plants; and (b) has already experienced a significant loss of biodiversity.

Two regions in North America meet these criteria: the California Floristic Province, a broad strip lying between the Sierra Nevada Range and the Pacific Coast and extending from southern Oregon to northern Mexico, contains about 3500 species of flowering plants of which approximately 61% are endemic; and the Madrean Pine–Oak Woodland features a series of unique habitats in the rugged mountain ranges and deep canyons in southern Arizona, southwestern Texas, and northern Mexico (Fig. 1.4). Unfortunately, only a small percentage of the land within each of these hotspots is currently protected, but national and international conservation

organizations, including The Nature Conservancy and Sierra Club, are actively working to protect additional habitats in both regions.

Patterns of distribution

Organisms are not randomly distributed but instead show more-or-less distinctive distributional patterns. Those with restricted distributions are said to be endemic to the area (e.g., pronghorns are endemic to western North America). Conversely, the patterns for some organisms are more complex and often difficult to explain. Discontinuous distributions, for example, are characterized by the presence of a single species in two or more widely separated areas. The range of wood ducks covers eastern North America, then skips the western plains and Rocky Mountains, but resumes along the northern Pacific coastline (California, Oregon, Washington, and southern British Columbia).

Continental patterns

The discontinuous distribution of some organisms involves even greater distances, which may even span parts of two or more continents. A few of the more common distributional patterns are briefly described below and, although they are traditionally applied to animals, they characterize plant distributions just as well.

Organisms whose distributions occur exclusively in North America are known as **Nearctic** species; prairie dogs and Gila monsters are good examples, as are pronghorns and wild turkeys. For comparison, **Palaearctic** species occur in Eurasia. Organisms found only in the Northern hemisphere – Nearctic and Palaearctic together – are said to have **Holarctic** distributions.

Holarctic distributions are most obvious in species associated with the boreal forest (e.g., lynx). Physical linkages between segments of boreal forest in North America and Eurasia are virtually absent, however, and thereby preclude movement of organisms between these regions. Lacking contact, these populations have varying degrees of reproductive isolation and progress towards speciation. Thus, Holarctic species are often distinguished by one or more subspecies (e.g., caribou and elk in the Nearctic and their Palaearctic counterparts, reindeer and red deer). In other cases, however, closely related species may represent Holarctic distributions (e.g., two species of beaver, one in North America and another in Europe).

Late in the 18th century, Holarctic species contributed to a transoceanic debate between Thomas Jefferson (1743–1826) and French naturalist Georges-Louis Buffon (1707–1788). Buffon, a scientific giant of his century, steadfastly maintained that the biota of North America was a degenerate and smaller counterpart of life in Europe. Jefferson, a champion of scientific rigor and America’s equality, responded by measuring specimens