



Stephen T. Thornton

Andrew Rex

Modern Physics

for Scientists and Engineers

Fourth Edition

This page intentionally left blank

Fundamental Constants

| Quantity | Symbol | Value(s) |
|--|--------------|---|
| Elementary charge | e | $1.6022 \times 10^{-19} \text{ C}$ |
| Speed of light in vacuum | c | $2.9979 \times 10^8 \text{ m/s}$ |
| Permeability of vacuum (magnetic constant) | μ_0 | $4\pi \times 10^{-7} \text{ N} \cdot \text{A}^{-2}$ |
| Permittivity of vacuum (electric constant) | ϵ_0 | $8.8542 \times 10^{-12} \text{ F} \cdot \text{m}^{-1}$ |
| Gravitation constant | G | $6.6738 \times 10^{-11} \text{ N} \cdot \text{m}^2 \cdot \text{kg}^{-2}$ |
| Planck constant | h | $6.6261 \times 10^{-34} \text{ J} \cdot \text{s}$ $4.1357 \times 10^{-15} \text{ eV} \cdot \text{s}$ |
| Avogadro constant | N_A | $6.0221 \times 10^{23} \text{ mol}^{-1}$ |
| Boltzmann constant | k | $1.3807 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$ |
| Stefan-Boltzmann constant | σ | $5.6704 \times 10^{-8} \text{ W} \cdot \text{m}^{-2} \cdot \text{K}^{-4}$ |
| Atomic mass unit | u | $1.66053886 \times 10^{-27} \text{ kg}$ $931.494061 \text{ MeV}/c^2$ |

Particle Masses

| Particle | Mass in units of | | |
|-------------------|--------------------------|--------------------|-------------------------|
| | kg | MeV/c ² | u |
| Electron | 9.1094×10^{-31} | 0.51100 | 5.4858×10^{-4} |
| Muon | 1.8835×10^{-28} | 105.66 | 0.11343 |
| Proton | 1.6726×10^{-27} | 938.27 | 1.00728 |
| Neutron | 1.6749×10^{-27} | 939.57 | 1.00866 |
| Deuteron | 3.3436×10^{-27} | 1875.61 | 2.01355 |
| α particle | 6.6447×10^{-27} | 3727.38 | 4.00151 |

Conversion Factors

| | |
|--|------------------------------------|
| 1 y = 3.156×10^7 s | 1 T = 10^4 G |
| 1 lightyear = 9.461×10^{15} m | 1 Ci = 3.7×10^{10} Bq |
| 1 cal = 4.186 J | 1 barn = 10^{-28} m ² |
| 1 MeV/c = 5.344×10^{-22} kg · m/s | 1 u = 1.66054×10^{-27} kg |
| 1 eV = 1.6022×10^{-19} J | |

Useful Combinations of Constants

$$\hbar = h/2\pi = 1.0546 \times 10^{-34} \text{ J} \cdot \text{s} = 6.5821 \times 10^{-16} \text{ eV} \cdot \text{s}$$

$$hc = 1.9864 \times 10^{-25} \text{ J} \cdot \text{m} = 1239.8 \text{ eV} \cdot \text{nm}$$

$$\hbar c = 3.1615 \times 10^{-26} \text{ J} \cdot \text{m} = 197.33 \text{ eV} \cdot \text{nm}$$

$$\frac{1}{4\pi\epsilon_0} = 8.9876 \times 10^9 \text{ N} \cdot \text{m}^2 \cdot \text{C}^{-2}$$

$$\text{Compton wavelength } \lambda_c = \frac{h}{m_e c} = 2.4263 \times 10^{-12} \text{ m}$$

$$\frac{e^2}{4\pi\epsilon_0} = 2.3071 \times 10^{-28} \text{ J} \cdot \text{m} = 1.4400 \times 10^{-9} \text{ eV} \cdot \text{m}$$

$$\text{Fine structure constant } \alpha = \frac{e^2}{4\pi\epsilon_0 \hbar c} = 0.0072974 \approx \frac{1}{137}$$

$$\text{Bohr magneton } \mu_B = \frac{e\hbar}{2m_e} = 9.2740 \times 10^{-24} \text{ J/T} = 5.7884 \times 10^{-5} \text{ eV/T}$$

$$\begin{aligned} \text{Nuclear magneton } \mu_N &= \frac{e\hbar}{2m_p} = 5.0508 \times 10^{-27} \text{ J/T} \\ &= 3.1525 \times 10^{-8} \text{ eV/T} \end{aligned}$$

$$\text{Bohr radius } a_0 = \frac{4\pi\epsilon_0 \hbar^2}{m_e e^2} = 5.2918 \times 10^{-11} \text{ m}$$

$$\text{Hydrogen ground state } E_0 = \frac{e^2}{8\pi\epsilon_0 a_0} = 13.606 \text{ eV} = 2.1799 \times 10^{-18} \text{ J}$$

$$\text{Rydberg constant } R_\infty = \frac{\alpha^2 m_e c}{2h} = 1.09737 \times 10^7 \text{ m}^{-1}$$

$$\text{Hydrogen Rydberg } R_H = \frac{\mu}{m_e} R_\infty = 1.09678 \times 10^7 \text{ m}^{-1}$$

$$\text{Gas constant } R = N_A k = 8.3145 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$$

$$\text{Magnetic flux quantum } \Phi_0 = \frac{h}{2e} = 2.0678 \times 10^{-15} \text{ T} \cdot \text{m}^2$$

$$\text{Classical electron radius } r_e = \alpha^2 a_0 = 2.8179 \times 10^{-15} \text{ m}$$

$$kT = 2.5249 \times 10^{-2} \text{ eV} \approx \frac{1}{40} \text{ eV at } T = 293 \text{ K}$$

Note: The latest values of the fundamental constants can be found at the National Institute of Standards and Technology website at <http://physics.nist.gov/cuu/Constants>



MODERN PHYSICS

For Scientists and Engineers

Fourth Edition

STEPHEN T. THORNTON

University of Virginia

ANDREW REX

University of Puget Sound



This page intentionally left blank

This is an electronic version of the print textbook. Due to electronic rights restrictions, some third party content may be suppressed. Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. The publisher reserves the right to remove content from this title at any time if subsequent rights restrictions require it. For valuable information on pricing, previous editions, changes to current editions, and alternate formats, please visit www.cengage.com/highered to search by ISBN#, author, title, or keyword for materials in your areas of interest.

Modern Physics for Scientists and Engineers,
Fourth Edition

Stephen T. Thornton, Andrew Rex

Publisher, Physics and Astronomy:
Charles Hartford

Development Editor: Ed Dodd

Associate Development Editor:
Brandi Kirksey

Editorial Assistant: Brendan Killion

Senior Media Editor: Rebecca Berardy
Schwartz

Marketing Manager: Jack Cooney

Marketing Coordinator: Julie Stefani

Marketing Communications Manager:
Darlene Macanan

Senior Content Project Manager:
Cathy Brooks

Senior Art Director: Cate Rickard Barr

Print Buyer: Diane Gibbons

Manufacturing Planner: Douglas Bertke

Rights Acquisitions Specialist: Shalice
Shah-Caldwell

Production Service and Compositor:
Graphic World Inc.

Text design: Reuter Designs

Cover design: Gary Regaglia, Metro Design

Cover Image: Eugene Kowaluk/University of
Rochester. In this experiment,
performed at the University of
Rochester's Laboratory for Laser
Energetics (LLE), 18 laser beams
delivered more than 5 kJ to a target
in 1-ns pulses.

© 2013, 2006, Brooks/Cole, Cengage Learning

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced, transmitted, stored, or used in any form or by any means, graphic, electronic, or mechanical, including but not limited to photocopying, recording, scanning, digitizing, taping, Web distribution, information networks, or information storage and retrieval systems, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the publisher.

For product information and technology assistance, contact us at
Cengage Learning Customer & Sales Support, 1-800-354-9706.

For permission to use material from this text or product,
submit all requests online at www.cengage.com/permissions.

Further permissions questions can be emailed to
permissionrequest@cengage.com.

Library of Congress Control Number: 2011926934

Student Edition:

ISBN-13: 978-1-133-10372-1

ISBN-10: 1-133-10372-3

Brooks/Cole

20 Channel Center Street
Boston, MA 02210
USA

Cengage Learning is a leading provider of customized learning solutions with office locations around the globe, including Singapore, the United Kingdom, Australia, Mexico, Brazil, and Japan. Locate your local office at
www.cengage.com/global.

Cengage Learning products are represented in Canada by Nelson Education, Ltd.

To learn more about Brooks/Cole, visit www.cengage.com/brookscole.

Purchase any of our products at your local college store or at our preferred online store www.CengageBrain.com

Cover image also used on pages: iii, iv, vi, xx; 1, 19, 33, 35, 38, 40, 41, 45, 51, 54, 57, 61, 63, 64, 67, 69, 71, 72, 84, 87, 90, 93, 94, 98, 101, 102, 106, 109, 110, 113, 116, 117, 120, 127, 129, 134, 136, 138, 139, 140, 146, 149, 150, 154, 156, 162, 167, 169, 170, 173, 174, 181, 184, 187, 188, 190, 195, 201, 203, 204, 205, 208, 211, 214, 215, 219, 222, 225, 228, 229, 230, 232, 234, 241, 246, 248, 249, 252, 254, 255, 257, 259, 260, 263, 265, 266, 272, 276, 280, 281, 285, 289, 290, 291, 293, 298, 303, 306, 310, 311, 315, 320, 322, 328, 330, 339, 342, 343, 348, 358, 364, 366, 374, 378, 382, 392, 399, 401, 402, 408, 412, 418, 431, 432, 436, 437, 438, 443, 444, 445, 447, 451, 452, 453, 458, 460, 461, 463, 467, 469, 475, 479, 480, 481, 484, 487, 488, 490, 496, 500, 502, 509, 512, 519, 522, 525, 531, 532, 533, 534, 535, 538, 544, 548, 550, 555, 562, 566, 567, 568, 577, 581, 583, 587, 588, 590, 591, 598, 602, 604, 610, A-1, A-2, A-4, A-6, A-7, A-9, A-12, A-14, A-37



Contents Overview

| | | |
|----|---|------|
| 1 | The Birth of Modern Physics | 1 |
| 2 | Special Theory of Relativity | 19 |
| 3 | The Experimental Basis of Quantum Physics | 84 |
| 4 | Structure of the Atom | 127 |
| 5 | Wave Properties of Matter and Quantum Mechanics I | 162 |
| 6 | Quantum Mechanics II | 201 |
| 7 | The Hydrogen Atom | 241 |
| 8 | Atomic Physics | 272 |
| 9 | Statistical Physics | 298 |
| 10 | Molecules, Lasers, and Solids | 339 |
| 11 | Semiconductor Theory and Devices | 392 |
| 12 | The Atomic Nucleus | 431 |
| 13 | Nuclear Interactions and Applications | 475 |
| 14 | Particle Physics | 519 |
| 15 | General Relativity | 555 |
| 16 | Cosmology and Modern Astrophysics— The Beginning and the End | 577 |
| | Appendices | A-1 |
| | Answers to Selected Odd-Numbered Problems | A-45 |
| | Index | I-1 |



Contents

Preface xii

Chapter 1

The Birth of Modern Physics 1

- 1.1 Classical Physics of the 1890s 2
 - Mechanics* 3
 - Electromagnetism* 4
 - Thermodynamics* 5
- 1.2 The Kinetic Theory of Gases 5
- 1.3 Waves and Particles 8
- 1.4 Conservation Laws and Fundamental Forces 10
 - Fundamental Forces* 10
- 1.5 The Atomic Theory of Matter 13
- 1.6 Unresolved Questions of 1895 and New Horizons 15
 - On the Horizon* 17
 - Summary 18

Chapter 2

Special Theory of Relativity 19

- 2.1 The Apparent Need for Ether 20
- 2.2 The Michelson-Morley Experiment 21

- 2.3 Einstein's Postulates 26
- 2.4 The Lorentz Transformation 29
- 2.5 Time Dilation and Length Contraction 31
 - Time Dilation* 31
 - Length Contraction* 35
- 2.6 Addition of Velocities 38
- 2.7 Experimental Verification 42
 - Muon Decay* 42
 - Atomic Clock Measurement* 43
 - Velocity Addition* 45
 - Testing Lorentz Symmetry* 46
- 2.8 Twin Paradox 46
- 2.9 Spacetime 48
- 2.10 Doppler Effect 52
 - Special Topic: Applications of the Doppler Effect* 54
- 2.11 Relativistic Momentum 58
- 2.12 Relativistic Energy 62
 - Total Energy and Rest Energy* 64
 - Equivalence of Mass and Energy* 65
 - Relationship of Energy and Momentum* 66
 - Massless Particles* 67
- 2.13 Computations in Modern Physics 68
 - Binding Energy* 70
- 2.14 Electromagnetism and Relativity 73
 - Summary 75

Chapter 3**The Experimental Basis of Quantum Physics 84**

- 3.1 Discovery of the X Ray and the Electron 84
- 3.2 Determination of Electron Charge 88
- 3.3 Line Spectra 91
Special Topic: The Discovery of Helium 93
- 3.4 Quantization 95
- 3.5 Blackbody Radiation 96
- 3.6 Photoelectric Effect 102
Experimental Results of Photoelectric Effect 103
Classical Interpretation 105
Einstein's Theory 107
Quantum Interpretation 107
- 3.7 X-Ray Production 110
- 3.8 Compton Effect 113
- 3.9 Pair Production and Annihilation 117
Summary 121

Chapter 4**Structure of the Atom 127**

- 4.1 The Atomic Models of Thomson and Rutherford 128
- 4.2 Rutherford Scattering 131
Special Topic: Lord Rutherford of Nelson 134
- 4.3 The Classical Atomic Model 139
- 4.4 The Bohr Model of the Hydrogen Atom 141
The Correspondence Principle 146
- 4.5 Successes and Failures of the Bohr Model 147
Reduced Mass Correction 148
Other Limitations 150
- 4.6 Characteristic X-Ray Spectra and Atomic Number 151
- 4.7 Atomic Excitation by Electrons 154
Summary 157

Chapter 5**Wave Properties of Matter and Quantum Mechanics I 162**

- 5.1 X-Ray Scattering 163
- 5.2 De Broglie Waves 168
Bohr's Quantization Condition 169
Special Topic: Cavendish Laboratory 170
- 5.3 Electron Scattering 172
- 5.4 Wave Motion 175
- 5.5 Waves or Particles? 182
- 5.6 Uncertainty Principle 186
- 5.7 Probability, Wave Functions, and the Copenhagen Interpretation 191
The Copenhagen Interpretation 192
- 5.8 Particle in a Box 194
Summary 196

Chapter 6**Quantum Mechanics II 201**

- 6.1 The Schrödinger Wave Equation 202
Normalization and Probability 204
Properties of Valid Wave Functions 206
Time-Independent Schrödinger Wave Equation 206
- 6.2 Expectation Values 209
- 6.3 Infinite Square-Well Potential 212
- 6.4 Finite Square-Well Potential 216
- 6.5 Three-Dimensional Infinite-Potential Well 218
- 6.6 Simple Harmonic Oscillator 220
- 6.7 Barriers and Tunneling 226
Potential Barrier with $E > V_0$ 226
Potential Barrier with $E < V_0$ 227
Potential Well 231
Alpha-Particle Decay 231
Special Topic: Scanning Probe Microscopes 232
Summary 235

Chapter 7**The Hydrogen Atom 241**

- 7.1 Application of the Schrödinger Equation to the Hydrogen Atom 241
- 7.2 Solution of the Schrödinger Equation for Hydrogen 242
Separation of Variables 243
Solution of the Radial Equation 244
Solution of the Angular and Azimuthal Equations 246
- 7.3 Quantum Numbers 248
Principal Quantum Number n 249
Orbital Angular Momentum Quantum Number ℓ 250
Magnetic Quantum Number m_ℓ 251
- 7.4 Magnetic Effects on Atomic Spectra—The Normal Zeeman Effect 253
- 7.5 Intrinsic Spin 258
Special Topic: Hydrogen and the 21-cm Line Transition 260
- 7.6 Energy Levels and Electron Probabilities 260
Selection Rules 262
Probability Distribution Functions 263
 Summary 268

Chapter 8**Atomic Physics 272**

- 8.1 Atomic Structure and the Periodic Table 273
Inert Gases 278
Alkalis 278
Alkaline Earths 278
Halogens 279
Transition Metals 279
Lanthanides 279
Special Topic: Rydberg Atoms 280
Actinides 281
- 8.2 Total Angular Momentum 281
Single-Electron Atoms 281
Many-Electron Atoms 285
LS Coupling 286
jj Coupling 289

- 8.3 Anomalous Zeeman Effect 292
 Summary 295

Chapter 9**Statistical Physics 298**

- 9.1 Historical Overview 299
- 9.2 Maxwell Velocity Distribution 301
- 9.3 Equipartition Theorem 303
- 9.4 Maxwell Speed Distribution 307
- 9.5 Classical and Quantum Statistics 311
Classical Distributions 311
Quantum Distributions 312
- 9.6 Fermi-Dirac Statistics 315
Introduction to Fermi-Dirac Theory 315
Classical Theory of Electrical Conduction 316
Quantum Theory of Electrical Conduction 317
- 9.7 Bose-Einstein Statistics 323
Blackbody Radiation 323
Liquid Helium 325
Special Topic: Superfluid ^3He 328
Symmetry of Boson Wave Functions 331
Bose-Einstein Condensation in Gases 332
 Summary 334

Chapter 10**Molecules, Lasers, and Solids 339**

- 10.1 Molecular Bonding and Spectra 340
Molecular Bonds 340
Rotational States 341
Vibrational States 342
Vibration and Rotation Combined 344
- 10.2 Stimulated Emission and Lasers 347
Scientific Applications of Lasers 352
Holography 353
Quantum Entanglement, Teleportation, and Information 354
Other Laser Applications 355
- 10.3 Structural Properties of Solids 356
- 10.4 Thermal and Magnetic Properties of Solids 359
Thermal Expansion 359
Thermal Conductivity 361

| | |
|---|-----|
| <i>Magnetic Properties</i> | 363 |
| <i>Diamagnetism</i> | 364 |
| <i>Paramagnetism</i> | 365 |
| <i>Ferromagnetism</i> | 366 |
| <i>Antiferromagnetism and Ferrimagnetism</i> | 367 |
| 10.5 Superconductivity | 367 |
| <i>The Search for a Higher T_c</i> | 374 |
| <i>Special Topic: Low-Temperature Methods</i> | 378 |
| <i>Other Classes of Superconductors</i> | 380 |
| 10.6 Applications of Superconductivity | 380 |
| <i>Josephson Junctions</i> | 381 |
| <i>Maglev</i> | 382 |
| <i>Generation and Transmission of Electricity</i> | 383 |
| <i>Other Scientific and Medical Applications</i> | 383 |
| Summary | 385 |

Chapter 11

Semiconductor Theory and Devices 392

| | |
|---|-----|
| 11.1 Band Theory of Solids | 392 |
| <i>Kronig-Penney Model</i> | 395 |
| <i>Band Theory and Conductivity</i> | 397 |
| 11.2 Semiconductor Theory | 397 |
| <i>Special Topic: The Quantum Hall Effect</i> | 402 |
| <i>Thermoelectric Effect</i> | 404 |
| 11.3 Semiconductor Devices | 406 |
| <i>Diodes</i> | 406 |
| <i>Rectifiers</i> | 408 |
| <i>Zener Diodes</i> | 408 |
| <i>Light-Emitting Diodes</i> | 409 |
| <i>Photovoltaic Cells</i> | 409 |
| <i>Transistors</i> | 413 |
| <i>Field Effect Transistors</i> | 415 |
| <i>Schottky Barriers</i> | 416 |
| <i>Semiconductor Lasers</i> | 417 |
| <i>Integrated Circuits</i> | 418 |
| 11.4 Nanotechnology | 421 |
| <i>Carbon Nanotubes</i> | 421 |
| <i>Nanoscale Electronics</i> | 422 |
| <i>Quantum Dots</i> | 424 |
| <i>Nanotechnology and the Life Sciences</i> | 425 |
| <i>Information Science</i> | 426 |
| Summary | 426 |

Chapter 12

The Atomic Nucleus 431

| | |
|--|-----|
| 12.1 Discovery of the Neutron | 431 |
| 12.2 Nuclear Properties | 434 |
| <i>Sizes and Shapes of Nuclei</i> | 435 |
| <i>Nuclear Density</i> | 437 |
| <i>Intrinsic Spin</i> | 437 |
| <i>Intrinsic Magnetic Moment</i> | 437 |
| <i>Nuclear Magnetic Resonance</i> | 438 |
| 12.3 The Deuteron | 439 |
| 12.4 Nuclear Forces | 441 |
| 12.5 Nuclear Stability | 442 |
| <i>Nuclear Models</i> | 448 |
| 12.6 Radioactive Decay | 449 |
| 12.7 Alpha, Beta, and Gamma Decay | 452 |
| <i>Alpha Decay</i> | 453 |
| <i>Beta Decay</i> | 456 |
| <i>Special Topic: Neutrino Detection</i> | 458 |
| <i>Gamma Decay</i> | 462 |
| 12.8 Radioactive Nuclides | 464 |
| <i>Time Dating Using Lead Isotopes</i> | 466 |
| <i>Radioactive Carbon Dating</i> | 467 |
| <i>Special Topic: The Formation and Age of the Earth</i> | 468 |
| Summary | 470 |

Chapter 13

Nuclear Interactions and Applications 475

| | |
|--------------------------------|-----|
| 13.1 Nuclear Reactions | 475 |
| <i>Cross Sections</i> | 478 |
| 13.2 Reaction Kinematics | 480 |
| 13.3 Reaction Mechanisms | 482 |
| <i>The Compound Nucleus</i> | 483 |
| <i>Direct Reactions</i> | 486 |
| 13.4 Fission | 486 |
| <i>Induced Fission</i> | 487 |
| <i>Thermal Neutron Fission</i> | 488 |
| <i>Chain Reactions</i> | 489 |

- 13.5 Fission Reactors 490
 - Nuclear Reactor Problems* 493
 - Breeder Reactors* 494
 - Future Nuclear Power Systems* 495
 - Special Topic: Early Fission Reactors* 496
- 13.6 Fusion 499
 - Formation of Elements* 499
 - Nuclear Fusion on Earth* 501
 - Controlled Thermonuclear Reactions* 502
- 13.7 Special Applications 505
 - Medicine* 505
 - Archaeology* 507
 - Art* 507
 - Crime Detection* 507
 - Mining and Oil* 508
 - Materials* 508
 - Small Power Systems* 510
 - New Elements* 510
 - Special Topic: The Search for New Elements* 512
- Summary 514

Chapter 14

Particle Physics 519

- 14.1 Early Discoveries 520
 - The Positron* 520
 - Yukawa's Meson* 521
- 14.2 The Fundamental Interactions 523
- 14.3 Classification of Particles 526
 - Leptons* 527
 - Hadrons* 528
 - Particles and Lifetimes* 530
- 14.4 Conservation Laws and Symmetries 532
 - Baryon Conservation* 532
 - Lepton Conservation* 533
 - Strangeness* 534
 - Symmetries* 535
- 14.5 Quarks 536
 - Quark Description of Particles* 537
 - Color* 539
 - Confinement* 539
- 14.6 The Families of Matter 541
- 14.7 Beyond the Standard Model 541
 - Neutrino Oscillations* 542
 - Matter-Antimatter* 542
 - Grand Unifying Theories* 543
 - Special Topic: Experimental Ingenuity* 544

- 14.8 Accelerators 546
 - Synchrotrons* 547
 - Linear Accelerators* 547
 - Fixed-Target Accelerators* 548
 - Colliders* 549
- Summary 551

Chapter 15

General Relativity 555

- 15.1 Tenets of General Relativity 555
 - Principle of Equivalence* 556
 - Spacetime Curvature* 558
- 15.2 Tests of General Relativity 560
 - Bending of Light* 560
 - Gravitational Redshift* 561
 - Perihelion Shift of Mercury* 562
 - Light Retardation* 563
- 15.3 Gravitational Waves 564
- 15.4 Black Holes 565
 - Special Topic: Gravitational Wave Detection* 566
- 15.5 Frame Dragging 572
- Summary 573

Chapter 16

Cosmology and Modern Astrophysics—The Beginning and the End 577

- 16.1 Evidence of the Big Bang 578
 - Hubble's Measurements* 578
 - Cosmic Microwave Background Radiation* 581
 - Nucleosynthesis* 581
 - Olbers' Paradox* 583
- 16.2 The Big Bang 583
- 16.3 Stellar Evolution 588
 - The Ultimate Fate of Stars* 589
 - Special Topic: Planck's Time, Length, and Mass* 591
- 16.4 Astronomical Objects 592
 - Active Galactic Nuclei and Quasars* 593
 - Gamma Ray Astrophysics* 594
 - Novae and Supernovae* 595

16.5 Problems with the Big Bang 599
The Inflationary Universe 599
The Lingering Problems 600

16.6 The Age of the Universe 603
Age of Chemical Elements 603
Age of Astronomical Objects 603
Cosmological Determinations 604
Universe Age Conclusion 607

16.7 The Standard Model of Cosmology 607

16.8 The Future 609
The Demise of the Sun 609
Special Topic: Future of Space Telescopes 610
The Future of the Universe 610
Are Other Earths Out There? 611
 Summary 612

Appendix 1
Fundamental Constants A-1

Appendix 2
Conversion Factors A-2

Appendix 3
Mathematical Relations A-4

Appendix 4
Periodic Table of the Elements A-6

Appendix 5
Mean Values and Distributions A-7

Appendix 6
Probability Integrals
 $I_n = \int_0^\infty x^n \exp(-ax^2) dx$ A-9

Appendix 7
Integrals of the Type $\int_0^\infty \frac{x^{n-1} dx}{e^x - 1}$ A-12

Appendix 8
Atomic Mass Table A-14

Appendix 9
Nobel Laureates in Physics A-37

Answers to Selected Odd-Numbered Problems A-45

Index I-1



Preface

Our objective in writing this book was to produce a textbook for a modern physics course of either one or two semesters for physics and engineering students. Such a course normally follows a full-year, introductory, calculus-based physics course for freshmen or sophomores. Before each edition we have the publisher send a questionnaire to users of modern physics books to see what needed to be changed or added. Most users like our textbook as is, especially the complete coverage of topics including the early quantum theory, subfields of physics, general relativity, and cosmology/astrophysics. Our book continues to be useful for either a one- or two-semester modern physics course. We have made no major changes in the order of subjects in the fourth edition.

Coverage

The first edition of our text established a trend for a contemporary approach to the exciting, thriving, and changing field of modern science. After briefly visiting the status of physics at the turn of the last century, we cover relativity and quantum theory, the basis of any study of modern physics. Almost all areas of science depend on quantum theory and the methods of experimental physics. We have included the name Quantum Mechanics in two of our chapter titles (Chapters 5 and 6) to emphasize the quantum connection. The latter part of the book is devoted to the subfields of physics (atomic, condensed matter, nuclear, and particle) and the exciting fields of cosmology and astrophysics. Our experience is that science and engineering majors particularly enjoy the study of modern physics after the sometimes-laborious study of classical mechanics, thermodynamics, electricity, magnetism, and optics. The level of mathematics is not difficult for the most part, and students feel they are finally getting to the frontiers of physics. We have brought the study of modern physics alive by presenting many current applications and challenges in physics, for example, nanoscience, high-temperature superconductors, quantum teleportation, neutrino mass and oscillations, missing dark mass and energy in the universe, gamma-ray bursts, holography, quantum dots, and nuclear fusion. Modern physics texts need to be updated periodically to include recent advances. Although we have emphasized modern applications, we also provide the sound theoretical basis for quantum theory that will be needed by physics majors in their upper division and graduate courses.

Changes for the Fourth Edition

Our book continues to be the most complete and up-to-date textbook in the modern physics market for sophomores/juniors. We have made several changes for the fourth edition to aid the student in learning modern physics. We have added additional end-of-chapter questions and problems and have modified many problems from earlier editions,

with an emphasis on including more real-world problems with current research applications whenever possible. We continue to have a larger number of questions and problems than competing textbooks, and for users of the robust Thornton/Rex *Modern Physics for Scientists and Engineers*, third edition course in WebAssign, we have a correlation guide of the fourth edition problems to that third edition course.

We have added additional examples to the already large number in the text. The pedagogical changes made for the third edition were highly successful. To encourage and support conceptual thinking by students, we continue to use conceptual examples and strategy discussion in the numerical examples. Examples with numerical solutions include a discussion of what needs to be accomplished in the example, the procedure to go through to find the answer, and relevant equations that will be needed. We present the example solutions in some detail, showing enough steps so that students can follow the solution to the end.

We continue to provide a significant number of photos and biographies of scientists who have made contributions to modern physics. We have done this to give students a perspective of the background, education, trials, and efforts of these scientists. We have also updated many of the Special Topic boxes, which we believe provide accurate and useful descriptions of the excitement of scientific discoveries, both past and current.

Chapter-by-Chapter Changes We have rewritten some sections in order to make the explanations clearer to the student. Some material has been deleted, and new material has been added. In particular we added new results that have been reported since the third edition. This is especially true for the chapters on the subfields of physics, Chapters 8–16. We have covered the most important subjects of modern physics, but we realize that in order to cover everything, the book would have to be much longer, which is not what our users want. Our intention is to keep the level of the textbook at the sophomore/junior undergraduate level. We think it is important for instructors to be able to supplement the book whenever they choose—especially to cover those topics in which they themselves are expert. Particular changes by chapter include the following:

- **Chapter 2:** we have updated the search for violations of Lorentz symmetry and added some discussion about four vectors.
- **Chapter 3:** we have rewritten the discussion of the Rayleigh-Jeans formula and Planck's discovery.
- **Chapter 9:** we improved the discussion about the symmetry of boson wave functions and its application to the Fermi exclusion principle and Bose-Einstein condensates.
- **Chapter 10:** we added a discussion of classes of superconductors and have updated our discussion concerning applications of superconductivity. The latter includes how superconductors are now being used to determine several fundamental constants.
- **Chapter 11:** we added more discussion about solar energy, Blu-ray DVD devices, increasing the number of transistors on a microchip using new semiconductor materials, graphene, and quantum dots. Our section on nanotechnology is especially complete.
- **Chapter 12:** we updated our discussion on neutrino detection and neutrino mass, added a description of nuclear magnetic resonance, and upgraded our discussion on using radioactive decay to study the oldest terrestrial materials.
- **Chapter 13:** we updated our discussion about nuclear power plants operating in the United States and the world and presented a discussion of possible new, improved reactors. We discussed the tsunami-induced tragedy at the Fukushima Daiichi nuclear power plant in Japan and added to our discussion of searches for new elements and their discoveries.
- **Chapter 14:** we upgraded our description of particle physics, improved and expanded the discussion on Feynman diagrams, updated the search for the Higgs boson, discussed new experiments on neutrino oscillations, and added discussion on matter-antimatter, supersymmetry, string theory, and M-theory. We mention that the LHC has begun operation as the Fermilab Tevatron accelerator is shutting down.
- **Chapter 15:** we improved our discussion on gravitational wave detection, added to our discussion on black holes, and included the final results of the Gravity Probe B satellite.

- **Chapter 16:** we changed the chapter name from Cosmology to Cosmology and Modern Astrophysics, because of the continued importance of the subject in modern physics. Our third edition of the textbook already had an excellent discussion and correct information about the age of the universe, dark matter, and dark energy, but Chapter 16 still has the most changes of any chapter, due to the current pace of research in the field. We have upgraded information and added discussion about Olbers' paradox, discovery of the cosmic microwave background, gamma ray astrophysics, standard model of cosmology, the future of space telescopes, and the future of the universe (Big Freeze, Big Crunch, Big Rip, Big Bounce, etc).

Teaching Suggestions

The text has been used extensively in its first three editions in courses at our home institutions. These include a one-semester course for physics and engineering students at the University of Virginia and a two-semester course for physics and pre-engineering students at the University of Puget Sound. These are representative of the one- and two-semester modern physics courses taught elsewhere. Both one- and two-semester courses should cover the material through the establishment of the periodic table in Chapter 8 with few exceptions. We have eliminated the denoting of optional sections, because we believe that depends on the wishes of the instructor, but we feel Sections 2.4, 4.2, 6.4, 6.6, 7.2, 7.6, 8.2, and 8.3 from the first nine chapters might be optional. Our suggestions for the one- and two-semester courses (3 or 4 credit hours per semester) are then

One-semester: Chapters 1–9 and selected other material as chosen by the instructor

Two-semester: Chapters 1–16 with supplementary material as desired, with possible student projects

An Internet-based, distance-learning version of the course is offered by one of the authors every summer (Physics 2620, 4 credit hours) at the University of Virginia that covers all chapters of the textbook, with emphasis on Chapters 1–8. Homework problems and exams are given on WebAssign. The course can be taken by a student located anywhere there is an Internet connection. See <http://modern.physics.virginia.edu/course/> for details.

Features

End-of-Chapter Problems

The 1166 questions and problems (258 questions and 908 problems) are more than in competing textbooks. Such a large number of questions and problems allows the instructor to make different homework assignments year after year without having to repeat problems. A correlation guide to the Thornton/Rex *Modern Physics for Scientists and Engineers*, third edition course in WebAssign is available via the Instructor's companion website (www.cengage.com/physics/thornton4e). We have tried to provide thought-provoking questions that have actual answers. In this edition we have focused on adding problems that have real-world or current research applications. The end-of-chapter problems have been separated by section, and general problems are included at the end to allow assimilation of the material. The easier problems are generally listed first within a section, and the more difficult ones are noted by a shaded blue square behind the problem number. A few computer-based problems are given in the text, but no computer disk supplement is provided, because many computer software programs are commercially available.

Solutions Manuals

PDF files of the *Instructor's Solutions Manual* are available to the instructor on the *Instructor's Resource CD-ROM* (by contacting your local Brooks/Cole—Cengage sales representative). This manual contains the *solution to every end-of-chapter problem* and has been checked by at

least two physics professors. The answers to selected odd-numbered problems are given at the end of the textbook itself. A *Student Solutions Manual* that contains the solutions to about 25% of the end-of-chapter problems is also available for sale to the students.

Instructor's Resource CD-ROM for Thornton/Rex's Modern Physics for Scientists and Engineers, Fourth Edition

Available to adopters is the *Modern Physics for Scientists and Engineers Instructor's Resource CD-ROM*. This CD-ROM includes PowerPoint® lecture outlines and also contains 200 pieces of line art from the text. It also features PDF files of the *Instructor's Solutions Manual*. Please guard this CD and do not let anyone have access to it. When end-of-chapter problem solutions find their way to the internet for sale, learning by students deteriorates because of the temptation to look up the solution.

Text Format

The two-color format helps to present clear illustrations and to highlight material in the text; for example, important and useful equations are highlighted in blue, and the most important part of each illustration is rendered in thick blue lines. Blue margin notes help guide the student to the important points, and the margins allow students to make their own notes. The first time key words or topics are introduced they are set in **boldface**, and *italics* are used for emphasis.

Examples

Although we had a large number of worked examples in the third edition, we have added new ones in this edition. The examples are written and presented in the manner in which students are expected to work the end-of-chapter problems: that is, to develop a conceptual understanding and strategy before attempting a numerical solution. Problem solving does not come easily for most students, especially the problems requiring several steps (that is, not simply plugging numbers into one equation). We expect that the many text examples with varying degrees of difficulty will help students.

Special Topic Boxes

Users have encouraged us to keep the Special Topic boxes. We believe both students and professors find them interesting, because they add some insight and detail into the excitement of physics. We have updated the material to keep them current.

History

We include historical aspects of modern physics that some students will find interesting and that others can simply ignore. We continue to include photos and biographies of scientists who have made significant contributions to modern physics. We believe this helps to enliven and humanize the material.

Website

Students can access the book's companion website at www.cengagebrain.com/shop/ISBN/9781133103721. This site features student study aids such as outlines, summaries, and conceptual questions for each chapter. Instructors will also find downloadable PowerPoint lectures and images for use in classroom lecture presentation. Students may also access the authors' websites at <http://www.modern.physics.virginia.edu/> and <http://www.pugetsound.edu/faculty-pages/rex> where the authors will post errata, present new exciting results, and give links to sites that have particularly interesting features like simulations and photos, among other things.

Acknowledgments

We acknowledge the assistance of many persons who have helped with this text. There are too many that helped us with the first three editions to list here, but the book would not have been possible without them. We acknowledge the professional staff at Brooks/Cole, Cengage Learning who helped make this fourth edition a useful, popular, and attractive text. They include Developmental Editor Ed Dodd and Senior Content Project Manager Cathy Brooks, who kept the production process on track, and Physics Publisher Charlie Hartford for his support, guidance, and encouragement. Elizabeth Budd did a superb job with the copyediting. We also want to thank Jeff Somers and the staff of Graphic World Inc. for their skilled efforts. We also want to thank the many individuals who gave us critical reviews and suggestions since the first edition. We especially would like to thank Michael Hood (Mt. San Antonio College) and Carol Hood (Augusta State University) for their help, especially with the Cosmology and Modern Astrophysics chapter. In preparing this fourth edition, we owe a special debt of gratitude to the following reviewers:

Jose D'Arruda, University of North Carolina, Pembroke
 David Church, Texas A & M University
 Hardin R. Dunham, Odessa College
 Paul A. Heiney, University of Pennsylvania
 Paul Keyes, Wayne State University
 Cody Martin, College of Menominee Nation

Prior to our work on this revision, we conducted a survey of professors to gauge how they taught their classes. In all, 78 professors responded with many insightful comments, and we would like to thank them for their feedback and suggestions.

We especially want to acknowledge the valuable help of Richard R. Bukrey of Loyola University of Chicago who helped us in many ways through his enlightening reviews, careful manuscript proofing, and checking of the end-of-chapter problem solutions in the first two editions, and to Thushara Perera of Illinois Wesleyan University, and Paul Weber of University of Puget Sound, for their accuracy review of the fourth edition. We also thank Allen Flora of Hood College for assuming the task of preparing and checking problem solutions for the third and fourth editions.

Stephen T. Thornton
 University of Virginia
 Charlottesville, Virginia
 stt@virginia.edu

Andrew Rex
 University of Puget Sound
 Tacoma, Washington
 rex@pugetsound.edu

The Birth of Modern Physics

1

CHAPTER

The more important fundamental laws and facts of physical science have all been discovered, and these are now so firmly established that the possibility of their ever being supplanted in consequence of new discoveries is exceedingly remote. . . . Our future discoveries must be looked for in the sixth place of decimals.

Albert A. Michelson, 1894

There is nothing new to be discovered in physics now. All that remains is more and more precise measurement.

William Thomson (Lord Kelvin), 1900

Although the Greek scholars Aristotle and Eratosthenes performed measurements and calculations that today we would call physics, the discipline of physics has its roots in the work of Galileo and Newton and others in the scientific revolution of the sixteenth and seventeenth centuries. The knowledge and practice of physics grew steadily for 200 to 300 years until another revolution in physics took place, which is the subject of this book. Physicists distinguish *classical physics*, which was mostly developed before 1895, from *modern physics*, which is based on discoveries made after 1895. The precise year is unimportant, but monumental changes occurred in physics around 1900.

The long reign of Queen Victoria of England, from 1837 to 1901, saw considerable changes in social, political, and intellectual realms, but perhaps none so important as the remarkable achievements that occurred in physics. For example, the description and predictions of electromagnetism by Maxwell are partly responsible for the rapid telecommunications of today. It was also during this period that thermodynamics rose to become an exact science. None of these achievements, however, have had the ramifications of the discoveries and applications of modern physics that would occur in the twentieth century. The world would never be the same.

In this chapter we briefly review the status of physics around 1895, including Newton's laws, Maxwell's equations, and the laws of thermodynamics. These results are just as important today as they were over a hundred years ago. Arguments by scientists concerning the interpretation of experimental data using

wave and particle descriptions that seemed to have been resolved 200 years ago were reopened in the twentieth century. Today we look back on the evidence of the late nineteenth century and wonder how anyone could have doubted the validity of the atomic view of matter. The fundamental interactions of gravity, electricity, and magnetism were thought to be well understood in 1895. Physicists continued to be driven by the goal of understanding fundamental laws throughout the twentieth century. This is demonstrated by the fact that other fundamental forces (specifically the nuclear and weak interactions) have been added, and in some cases—curious as it may seem—various forces have even been combined. The search for the holy grail of fundamental interactions continues unabated today.

We finish this chapter with a status report on physics just before 1900. The few problems not then understood would be the basis for decades of fruitful investigations and discoveries continuing into the twenty-first century. We hope you find this chapter interesting both for the physics presented and for the historical account of some of the most exciting scientific discoveries of the modern era.

1.1 Classical Physics of the 1890s

Scientists and engineers of the late nineteenth century were indeed rather smug. They thought they had just about everything under control (see the quotes from Michelson and Kelvin on page 1). The best scientists of the day were highly recognized and rewarded. Public lectures were frequent. Some scientists had easy access to their political leaders, partly because science and engineering had benefited their war machines, but also because of the many useful technical advances. Basic research was recognized as important because of the commercial and military applications of scientific discoveries. Although there were only primitive automobiles and no airplanes in 1895, advances in these modes of transportation were soon to follow. A few people already had telephones, and plans for widespread distribution of electricity were under way.

Based on their success with what we now call macroscopic classical results, scientists felt that given enough time and resources, they could explain just about anything. They did recognize some difficult questions they still couldn't answer; for example, they didn't clearly understand the structure of matter—that was under intensive investigation. Nevertheless, on a macroscopic scale, they knew how to build efficient engines. Ships plied the lakes, seas, and oceans of the world. Travel between the countries of Europe was frequent and easy by train. Many scientists were born in one country, educated in one or two others, and eventually worked in still other countries. The most recent ideas traveled relatively quickly among the centers of research. Except for some isolated scientists, of whom Einstein is the most notable example, discoveries were quickly and easily shared. Scientific journals were becoming accessible.

The ideas of classical physics are just as important and useful today as they were at the end of the nineteenth century. For example, they allow us to build automobiles and produce electricity. The conservation laws of energy, linear momentum, angular momentum, and charge can be stated as follows:

Early successes of science

Classical conservation laws

Conservation of energy: The total sum of energy (in all its forms) is conserved in all interactions.

Conservation of linear momentum: In the absence of external forces, linear momentum is conserved in all interactions (vector relation).

Conservation of angular momentum: In the absence of external torque, angular momentum is conserved in all interactions (vector relation).

Conservation of charge: Electric charge is conserved in all interactions.

A nineteenth-century scientist might have added the **conservation of mass** to this list, but we know it not to be valid today (you will find out why in Chapter 2). These conservation laws are reflected in the laws of mechanics, electromagnetism, and thermodynamics. Electricity and magnetism, separate subjects for hundreds of years, were combined by James Clerk Maxwell (1831–1879) in his four equations. Maxwell showed optics to be a special case of electromagnetism. Waves, which permeated mechanics and optics, were known to be an important component of nature. Many natural phenomena could be explained by wave motion using the laws of physics.

Mechanics

The laws of mechanics were developed over hundreds of years by many researchers. Important contributions were made by astronomers because of the great interest in the heavenly bodies. Galileo (1564–1642) may rightfully be called the first great experimenter. His experiments and observations laid the groundwork for the important discoveries to follow during the next 200 years.

Isaac Newton (1642–1727) was certainly the greatest scientist of his time and one of the best the world has ever seen. His discoveries were in the fields of mathematics, astronomy, and physics and include gravitation, optics, motion, and forces.

We owe to Newton our present understanding of motion. He understood clearly the relationships among position, displacement, velocity, and acceleration. He understood how motion was possible and that a body at rest was just a special case of a body having constant velocity. It may not be so apparent to us today, but we should not forget the tremendous unification that Newton made when he pointed out that the motions of the planets about our sun can be understood by the same laws that explain motion on Earth, like apples falling from trees or a soccer ball being shot toward a goal. Newton was able to elucidate

Galileo, the first great experimenter

Newton, the greatest scientist of his time



Scala/Art Resource, NY

Galileo Galilei (1564–1642) was born, educated, and worked in Italy. Often said to be the “father of physics” because of his careful experimentation, he is shown here performing experiments by rolling balls on an inclined plane. He is perhaps best known for his experiments on motion, the development of the telescope, and his many astronomical discoveries. He came into disfavor with the Catholic Church for his belief in the Copernican theory. He was finally cleared of heresy by Pope John Paul II in 1992, 350 years after his death.

Newton's laws



Courtesy of Bausch & Lomb Optical Co. and the AIP Niels Bohr Library.

Isaac Newton (1642–1727), the great English physicist and mathematician, did most of his work at Cambridge where he was educated and became the Lucasian Professor of Mathematics. He was known not only for his work on the laws of motion but also as a founder of optics. His useful works are too numerous to list here, but it should be mentioned that he spent a considerable amount of his time on alchemy, theology, and the spiritual universe. His writings on these subjects, which were dear to him, were quite unorthodox. This painting shows him performing experiments with light.

Maxwell's equations

carefully the relationship between net force and acceleration, and his concepts were stated in three laws that bear his name today:

Newton's first law: *An object in motion with a constant velocity will continue in motion unless acted upon by some net external force.* A body at rest is just a special case of Newton's first law with zero velocity. Newton's first law is often called the *law of inertia* and is also used to describe inertial reference frames.

Newton's second law: *The acceleration \vec{a} of a body is proportional to the net external force \vec{F} and inversely proportional to the mass m of the body. It is stated mathematically as*

$$\vec{F} = m\vec{a} \quad (1.1a)$$

A more general statement* relates force to the time rate of change of the linear momentum \vec{p} .

$$\vec{F} = \frac{d\vec{p}}{dt} \quad (1.1b)$$

Newton's third law: *The force exerted by body 1 on body 2 is equal in magnitude and opposite in direction to the force that body 2 exerts on body 1.* If the force on body 2 by body 1 is denoted by \vec{F}_{21} , then Newton's third law is written as

$$\vec{F}_{21} = -\vec{F}_{12} \quad (1.2)$$

It is often called the *law of action and reaction*.

These three laws develop the concept of force. Using that concept together with the concepts of velocity \vec{v} , acceleration \vec{a} , linear momentum \vec{p} , rotation (angular velocity $\vec{\omega}$ and angular acceleration $\vec{\alpha}$), and angular momentum \vec{L} , we can describe the complex motion of bodies.

Electromagnetism

Electromagnetism developed over a long period of time. Important contributions were made by Charles Coulomb (1736–1806), Hans Christian Oersted (1777–1851), Thomas Young (1773–1829), André Ampère (1775–1836), Michael Faraday (1791–1867), Joseph Henry (1797–1878), James Clerk Maxwell (1831–1879), and Heinrich Hertz (1857–1894). Maxwell showed that electricity and magnetism were intimately connected and were related by a change in the inertial frame of reference. His work also led to the understanding of electromagnetic radiation, of which light and optics are special cases. Maxwell's four equations, together with the Lorentz force law, explain much of electromagnetism.

$$\text{Gauss's law for electricity} \quad \oint \vec{E} \cdot d\vec{A} = \frac{q}{\epsilon_0} \quad (1.3)$$

$$\text{Gauss's law for magnetism} \quad \oint \vec{B} \cdot d\vec{A} = 0 \quad (1.4)$$

$$\text{Faraday's law} \quad \oint \vec{E} \cdot d\vec{s} = -\frac{d\Phi_B}{dt} \quad (1.5)$$

*It is a remarkable fact that Newton wrote his second law not as $\vec{F} = m\vec{a}$, but as $\vec{F} = d(m\vec{v})/dt$, thus taking into account mass flow and change in velocity. This has applications in both fluid mechanics and rocket propulsion.

$$\text{Generalized Ampere's law} \quad \oint \vec{B} \cdot d\vec{s} = \mu_0 \epsilon_0 \frac{d\Phi_E}{dt} + \mu_0 I \quad (1.6)$$

$$\text{Lorentz force law} \quad \vec{F} = q\vec{E} + q\vec{v} \times \vec{B} \quad (1.7)$$

Maxwell's equations indicate that charges and currents create fields, and in turn, these fields can create other fields, both electric and magnetic.

Thermodynamics

Thermodynamics deals with temperature T , heat Q , work W , and the internal energy of systems U . The understanding of the concepts used in thermodynamics—such as pressure P , volume V , temperature, thermal equilibrium, heat, entropy, and especially energy—was slow in coming. We can understand the concepts of pressure and volume as mechanical properties, but the concept of temperature must be carefully considered. We have learned that the internal energy of a system of noninteracting point masses depends only on the temperature.

Important contributions to thermodynamics were made by Benjamin Thompson (Count Rumford, 1753–1814), Sadi Carnot (1796–1832), James Joule (1818–1889), Rudolf Clausius (1822–1888), and William Thomson (Lord Kelvin, 1824–1907). The primary results of thermodynamics can be described in two laws:

First law of thermodynamics: *The change in the internal energy ΔU of a system is equal to the heat Q added to the system plus the work W done on the system.*

$$\Delta U = Q + W \quad (1.8)$$

The first law of thermodynamics generalizes the conservation of energy by including heat.

Second law of thermodynamics: *It is not possible to convert heat completely into work without some other change taking place.* Various forms of the second law state similar, but slightly different, results. For example, it is not possible to build a perfect engine or a perfect refrigerator. It is not possible to build a perpetual motion machine. Heat does not spontaneously flow from a colder body to a hotter body without some other change taking place. The second law forbids all these from happening. The first law states the conservation of energy, but the second law says what kinds of energy processes cannot take place. For example, it is possible to completely convert work into heat, but not vice versa, without some other change taking place.

Two other “laws” of thermodynamics are sometimes expressed. One is called the “zeroth” law, and it is useful in understanding temperature. It states that *if two thermal systems are in thermodynamic equilibrium with a third system, they are in equilibrium with each other.* We can state it more simply by saying that *two systems at the same temperature as a third system have the same temperature as each other.* This concept was not explicitly stated until the twentieth century. The “third” law of thermodynamics expresses that *it is not possible to achieve an absolute zero temperature.*

1.2 The Kinetic Theory of Gases

We understand now that gases are composed of atoms and molecules in rapid motion, bouncing off each other and the walls, but in the 1890s this had just gained acceptance. The kinetic theory of gases is related to thermodynamics and

Laws of thermodynamics

to the atomic theory of matter, which we discuss in Section 1.5. Experiments were relatively easy to perform on gases, and the Irish chemist Robert Boyle (1627–1691) showed around 1662 that the pressure times the volume of a gas was constant for a constant temperature. The relation $PV = \text{constant}$ (for constant T) is now referred to as *Boyle's law*. The French physicist Jacques Charles (1746–1823) found that $V/T = \text{constant}$ (at constant pressure), referred to as *Charles's law*. Joseph Louis Gay-Lussac (1778–1850) later produced the same result, and the law is sometimes associated with his name. If we combine these two laws, we obtain the ideal gas equation

Ideal gas equation

$$PV = nRT \quad (1.9)$$

where n is the number of moles and R is the ideal gas constant, $8.31 \text{ J/mol} \cdot \text{K}$.

In 1811 the Italian physicist Amedeo Avogadro (1776–1856) proposed that equal volumes of gases at the same temperature and pressure contained equal numbers of molecules. This hypothesis was so far ahead of its time that it was not accepted for many years. The famous English chemist John Dalton opposed the idea because he apparently misunderstood the difference between atoms and molecules. Considering the rudimentary nature of the atomic theory of matter at the time, this was not surprising.

Daniel Bernoulli (1700–1782) apparently originated the kinetic theory of gases in 1738, but his results were generally ignored. Many scientists, including Newton, Laplace, Davy, Herapath, and Waterston, had contributed to the development of kinetic theory by 1850. Theoretical calculations were being compared with experiments, and by 1895 the kinetic theory of gases was widely accepted. The statistical interpretation of thermodynamics was made in the latter half of the nineteenth century by Maxwell, the Austrian physicist Ludwig Boltzmann (1844–1906), and the American physicist J. Willard Gibbs (1839–1903).

In introductory physics classes, the kinetic theory of gases is usually taught by applying Newton's laws to the collisions that a molecule makes with other molecules and with the walls. A representation of a few molecules colliding is shown in Figure 1.1. In the simple model of an ideal gas, only elastic collisions are considered. By taking averages over the collisions of many molecules, the ideal gas law, Equation (1.9), is revealed. The average kinetic energy of the molecules is shown to be linearly proportional to the temperature, and the internal energy U is

$$U = nN_A \langle K \rangle = \frac{3}{2} nRT \quad (1.10)$$

where n is the number of moles of gas, N_A is Avogadro's number, $\langle K \rangle$ is the average kinetic energy of a molecule, and R is the ideal gas constant. This relation ignores any nontranslational contributions to the molecular energy, such as rotations and vibrations.

However, energy is not represented only by translational motion. It became clear that all *degrees of freedom*, including rotational and vibrational, were also capable of carrying energy. The *equipartition theorem* states that each degree of freedom of a molecule has an average energy of $kT/2$, where k is the Boltzmann constant ($k = R/N_A$). Translational motion has three degrees of freedom, and rotational and vibrational modes can also be excited at higher temperatures. If there are f degrees of freedom, then Equation (1.10) becomes

$$U = \frac{f}{2} nRT \quad (1.11)$$




Figure 1.1 Molecules inside a closed container are shown colliding with the walls and with each other. The motions of a few molecules are indicated by the arrows. The number of molecules inside the container is huge.

Statistical thermodynamics

Equipartition theorem

Internal energy

The molar ($n = 1$) heat capacity c_V at constant volume for an ideal gas is the rate of change in internal energy with respect to change in temperature and is given by

$$c_V = \frac{3}{2}R \quad (1.12) \quad \text{Heat capacity}$$

The experimental quantity c_V/R is plotted versus temperature for hydrogen in Figure 1.2. The ratio c_V/R is equal to $3/2$ for low temperatures, where only translational kinetic energy is important, but it rises to $5/2$ at 300 K, where rotations occur for H_2 , and finally reaches $7/2$, because of vibrations at still higher temperatures, before the molecule dissociates. Although the kinetic theory of gases fails to predict specific heats for real gases, it leads to models that can be used on a gas-by-gas basis. Kinetic theory is also able to provide useful information on other properties such as diffusion, speed of sound, mean free path, and collision frequency.

In the 1850s Maxwell derived a relation for the distribution of speeds of the molecules in gases. The distribution of speeds $f(v)$ is given as a function of the speed and the temperature by the equation

$$f(v) = 4\pi N \left(\frac{m}{2\pi kT} \right)^{3/2} v^2 e^{-mv^2/2kT} \quad (1.13) \quad \text{Maxwell's speed distribution}$$

where m is the mass of a molecule and T is the temperature. This result is plotted for nitrogen in Figure 1.3 for temperatures of 300 K, 1000 K, and 4000 K. The peak of each distribution is the most probable speed of a gas molecule for the given temperature. In 1895 measurement was not precise enough to confirm Maxwell's distribution, and it was not confirmed experimentally until 1921.

By 1895 Boltzmann had made Maxwell's calculation more rigorous, and the general relation is called the *Maxwell-Boltzmann distribution*. The distribution can be used to find the *root-mean-square* speed v_{rms} ,

$$v_{\text{rms}} = \sqrt{\langle v^2 \rangle} = \sqrt{\frac{3kT}{m}} \quad (1.14)$$

which shows the relationship of the energy to the temperature for an ideal gas:

$$U = nN_A \langle K \rangle = nN_A \frac{m\langle v^2 \rangle}{2} = nN_A \frac{m3kT}{2m} = \frac{3}{2} nRT \quad (1.15)$$

This was the result of Equation (1.10).





Figure 1.2 The molar heat capacity at constant volume (c_V) divided by R (c_V/R is dimensionless) is displayed as a function of temperature for hydrogen gas. Note that as the temperature increases, the rotational and vibrational modes become important. This experimental result is consistent with the equipartition theorem, which adds $kT/2$ of energy per molecule ($RT/2$ per mole) for each degree of freedom.

Figure 1.3 The Maxwell distribution of molecular speeds (for nitrogen), $f(v)$, is shown as a function of speed for three temperatures.



1.3 Waves and Particles

We first learned the concepts of velocity, acceleration, force, momentum, and energy in introductory physics by using a single particle with its mass concentrated in one small point. In order to adequately describe nature, we add two- and three-dimensional bodies and rotations and vibrations. However, many aspects of physics can still be treated as if the bodies are simple particles. In particular, the kinetic energy of a moving particle is one way that energy can be transported from one place to another.

Energy transport

But we have found that many natural phenomena can be explained only in terms of *waves*, which are traveling disturbances that carry energy. This description includes standing waves, which are superpositions of traveling waves. Most waves, like water waves and sound waves, need an elastic medium in which to move. Curiously enough, matter is not transported in waves—but energy is. Mass may oscillate, but it doesn't actually propagate along with the wave. Two examples are a cork and a boat on water. As a water wave passes, the cork gains energy as it moves up and down, and after the wave passes, the cork remains. The boat also reacts to the wave, but it primarily rocks back and forth, throwing around things that are not fixed on the boat. The boat obtains considerable kinetic energy from the wave. After the wave passes, the boat eventually returns to rest.

Nature of light: waves or particles?

Waves and particles were the subject of disagreement as early as the seventeenth century, when there were two competing theories of the nature of light. Newton supported the idea that light consisted of corpuscles (or particles). He performed extensive experiments on light for many years and finally published his book *Opticks* in 1704. *Geometrical optics* uses straight-line, particle-like trajectories called *rays* to explain familiar phenomena such as reflection and refraction. Geometrical optics was also able to explain the apparent observation of sharp shadows. The competing theory considered light as a wave phenomenon. Its strongest proponent was the Dutch physicist Christian Huygens (1629–1695), who presented his theory in 1678. The wave theory could also explain reflection and refraction, but it could not explain the sharp shadows observed. Experimental physics of the 1600s and 1700s was not able to discern between the two competing theories. Huygens's poor health and other duties kept him from working on optics much after 1678. Although Newton did not feel strongly about his corpuscular

theory, the magnitude of his reputation caused it to be almost universally accepted for more than a hundred years and throughout most of the eighteenth century.

Finally, in 1802, the English physician Thomas Young (1773–1829) announced the results of his two-slit interference experiment, indicating that light behaved as a wave. Even after this singular event, the corpuscular theory had its supporters. During the next few years Young and, independently, Augustin Fresnel (1788–1827) performed several experiments that clearly showed that light behaved as a wave. By 1830 most physicists believed in the wave theory—some 150 years after Newton performed his first experiments on light.

One final experiment indicated that the corpuscular theory was difficult to accept. Let c be the speed of light in vacuum and v be the speed of light in another medium. If light behaves as a particle, then to explain refraction, light must speed up when going through denser material ($v > c$). The wave theory of Huygens predicts just the opposite ($v < c$). The measurements of the speed of light in various media were slowly improving, and finally, in 1850, Foucault showed that *light traveled more slowly in water than in air*. The corpuscular theory seemed incorrect. Newton would probably have been surprised that his weakly held beliefs lasted as long as they did. Now we realize that geometrical optics is correct only if the wavelength of light is much smaller than the size of the obstacles and apertures that the light encounters.

Figure 1.4 shows the “shadows” or *diffraction patterns* from light falling on sharp edges. In Figure 1.4a the alternating black and white lines can be seen all around the razor blade’s edges. Figure 1.4b is a highly magnified photo of the diffraction from a sharp edge. The bright and dark regions can be understood only if light is a wave and not a particle. The physicists of 200 to 300 years ago apparently did not observe such phenomena. They believed that shadows were sharp, and only the particle nature of light could explain their observations.

In the 1860s Maxwell showed that electromagnetic waves consist of oscillating electric and magnetic fields. Visible light covers just a narrow range of the total electromagnetic spectrum, and all electromagnetic radiation travels at the speed of light c in free space, given by

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}} = \lambda f \quad (1.16)$$

where λ is the wavelength and f is the frequency. The fundamental constants μ_0 and ϵ_0 are defined in electricity and magnetism and reveal the connection to the speed of light. In 1887 the German physicist Heinrich Hertz (1857–1894) succeeded in generating and detecting electromagnetic waves having wavelengths




Figure 1.4 In contradiction to what scientists thought in the seventeenth century, shadows are not sharp, but show dramatic diffraction patterns—as seen here (a) for a razor blade and (b) for a highly magnified sharp edge.