

CHAPTER 1

CHEMICAL FOUNDATIONS

Questions

19. A law summarizes what happens, e.g., law of conservation of mass in a chemical reaction or the ideal gas law, $PV = nRT$. A theory (model) is an attempt to explain why something happens. Dalton's atomic theory explains why mass is conserved in a chemical reaction. The kinetic molecular theory explains why pressure and volume are inversely related at constant temperature and moles of gas present, as well as explaining the other mathematical relationships summarized in $PV = nRT$.
20. A dynamic process is one that is active as opposed to static. In terms of the scientific method, scientists are always performing experiments to prove or disprove a hypothesis or a law or a theory. Scientists do not stop asking questions just because a given theory seems to account satisfactorily for some aspect of natural behavior. The key to the scientific method is to continually ask questions and perform experiments. Science is an active process, not a static one.
21. The fundamental steps are
- (1) making observations;
 - (2) formulating hypotheses;
 - (3) performing experiments to test the hypotheses.

The key to the scientific method is performing experiments to test hypotheses. If after the test of time the hypotheses seem to account satisfactorily for some aspect of natural behavior, then the set of tested hypotheses turns into a theory (model). However, scientists continue to perform experiments to refine or replace existing theories.

22. A random error has equal probability of being too high or too low. This type of error occurs when estimating the value of the last digit of a measurement. A systematic error is one that always occurs in the same direction, either too high or too low. For example, this type of error would occur if the balance you were using weighed all objects 0.20 g too high, that is, if the balance wasn't calibrated correctly. A random error is an indeterminate error, whereas a systematic error is a determinate error.
23. A qualitative observation expresses what makes something what it is; it does not involve a number; e.g., the air we breathe is a mixture of gases, ice is less dense than water, rotten milk stinks.

The SI units are mass in kilograms, length in meters, and volume in the derived units of m^3 . The assumed uncertainty in a number is ± 1 in the last significant figure of the number. The precision of an instrument is related to the number of significant figures associated with an

experimental reading on that instrument. Different instruments for measuring mass, length, or volume have varying degrees of precision. Some instruments only give a few significant figures for a measurement, whereas others will give more significant figures.

24. Precision: reproducibility; accuracy: the agreement of a measurement with the true value.
- Imprecise and inaccurate data: 12.32 cm, 9.63 cm, 11.98 cm, 13.34 cm
 - Precise but inaccurate data: 8.76 cm, 8.79 cm, 8.72 cm, 8.75 cm
 - Precise and accurate data: 10.60 cm, 10.65 cm, 10.63 cm, 10.64 cm

Data can be imprecise if the measuring device is imprecise as well as if the user of the measuring device has poor skills. Data can be inaccurate due to a systematic error in the measuring device or with the user. For example, a balance may read all masses as weighing 0.2500 g too high or the user of a graduated cylinder may read all measurements 0.05 mL too low.

A set of measurements that are imprecise implies that all the numbers are not close to each other. If the numbers aren't reproducible, then all the numbers can't be very close to the true value. Some say that if the average of imprecise data gives the true value, then the data are accurate; a better description is that the data takers are extremely lucky.

25. Significant figures are the digits we associate with a number. They contain all of the certain digits and the first uncertain digit (the first estimated digit). What follows is one thousand indicated to varying numbers of significant figures: 1000 or 1×10^3 (1 S.F.); 1.0×10^3 (2 S.F.); 1.00×10^3 (3 S.F.); 1000. or 1.000×10^3 (4 S.F.).

To perform the calculation, the addition/subtraction significant figure rule is applied to $1.5 - 1.0$. The result of this is the one-significant-figure answer of 0.5. Next, the multiplication/division rule is applied to $0.5/0.50$. A one-significant-figure number divided by a two-significant-figure number yields an answer with one significant figure (answer = 1).

26. From Figure 1.9 of the text, a change in temperature of 180°F is equal to a change in temperature of 100°C and 100 K. A degree unit on the Fahrenheit scale is not as large as a degree unit on the Celsius or Kelvin scales. Therefore, a 20° change in the Celsius or Kelvin temperature would correspond to a larger temperature change than a 20° change in the Fahrenheit scale. The 20° temperature change on the Celsius and Kelvin scales are equal to each other.
27. Straight line equation: $y = mx + b$, where m is the slope of the line and b is the y-intercept. For the T_F vs. T_C plot:

$$T_F = (9/5)T_C + 32$$
$$y = m x + b$$

The slope of the plot is 1.8 (= 9/5) and the y-intercept is 32°F .

For the T_C vs. T_K plot:

$$T_C = T_K - 273$$
$$y = m x + b$$

The slope of the plot is 1, and the y-intercept is -273°C .

34. a. 100; 1 S.F. b. 1.0×10^2 ; 2 S.F. c. 1.00×10^3 ; 3 S.F.
 d. 100.; 3 S.F. e. 0.0048; 2 S.F. f. 0.00480; 3 S.F.
 g. 4.80×10^{-3} ; 3 S.F. h. 4.800×10^{-3} ; 4 S.F.

35. When rounding, the last significant figure stays the same if the number after this significant figure is less than 5 and increases by one if the number is greater than or equal to 5.

- a. 3.42×10^{-4} b. 1.034×10^4 c. 1.7992×10^1 d. 3.37×10^5
 36. a. 4×10^5 b. 3.9×10^5 c. 3.86×10^5 d. 3.8550×10^5

37. Volume measurements are estimated to one place past the markings on the glassware. The first graduated cylinder is labeled to 0.2 mL volume increments, so we estimate volumes to the hundredths place. Realistically, the uncertainty in this graduated cylinder is ± 0.05 mL. The second cylinder, with 0.02 mL volume increments, will have an uncertainty of ± 0.005 mL. The approximate volume in the first graduated cylinder is 2.85 mL, and the volume in the other graduated cylinder is approximately 0.280 mL. The total volume would be:

$$\begin{array}{r} 2.85 \text{ mL} \\ +0.280 \text{ mL} \\ \hline 3.13 \text{ mL} \end{array}$$

We should report the total volume to the hundredths place because the volume from the first graduated cylinder is only read to the hundredths (read to two decimal places). The first graduated cylinder is the least precise volume measurement because the uncertainty of this instrument is in the hundredths place, while the uncertainty of the second graduated cylinder is to the thousandths place. It is always the least precise measurement that limits the precision of a calculation.

38. a. Volumes are always estimated to one position past the marked volume increments. The estimated volume of the first beaker is 32.7 mL, the estimated volume of the middle beaker is 33 mL, and the estimated volume in the last beaker is 32.73 mL.
 b. Yes, all volumes could be identical to each other because the more precise volume readings can be rounded to the other volume readings. But because the volumes are in three different measuring devices, each with its own unique uncertainty, we cannot say with certainty that all three beakers contain the same amount of water.
 c. 32.7 mL
 33 mL
 32.73 mL
 98.43 mL = 98 mL

The volume in the middle beaker can only be estimated to the ones place, which dictates that the sum of the volume should be reported to the ones place. As is always the case, the least precise measurement determines the precision of a calculation.

39. For addition and/or subtraction, the result has the same number of decimal places as the number in the calculation with the fewest decimal places. When the result is rounded to the correct number of significant figures, the last significant figure stays the same if the number

after this significant figure is less than 5 and increases by one if the number is greater than or equal to 5. The underline shows the last significant figure in the intermediate answers.

- a. $212.2 + 26.7 + 402.09 = 640.\underline{99} = 641.0$
- b. $1.0028 + 0.221 + 0.10337 = 1.32\underline{7}17 = 1.327$
- c. $52.331 + 26.01 - 0.9981 = 77.34\underline{29} = 77.34$
- d. $2.01 \times 10^2 + 3.014 \times 10^3 = 2.01 \times 10^2 + 30.14 \times 10^2 = 32.1\underline{5} \times 10^2 = 3215$

When the exponents are different, it is easiest to apply the addition/subtraction rule when all numbers are based on the same power of 10.

- e. $7.255 - 6.8350 = 0.42 = 0.420$ (first uncertain digit is in the third decimal place).

40. For multiplication and/or division, the result has the same number of significant figures as the number in the calculation with the fewest significant figures.

- a. $\frac{0.102 \times 0.0821 \times 273}{1.01} = \underline{2.2635} = 2.26$
- b. $0.14 \times 6.022 \times 10^{23} = \underline{8.431} \times 10^{22} = 8.4 \times 10^{22}$; since 0.14 only has two significant figures, the result should only have two significant figures.
- c. $4.0 \times 10^4 \times 5.021 \times 10^{-3} \times 7.34993 \times 10^2 = \underline{1.476} \times 10^5 = 1.5 \times 10^5$
- d. $\frac{2.00 \times 10^6}{3.00 \times 10^{-7}} = \underline{6.6667} \times 10^{12} = 6.67 \times 10^{12}$

41. a. For this problem, apply the multiplication/division rule first; then apply the addition/subtraction rule to arrive at the one-decimal-place answer. We will generally round off at intermediate steps in order to show the correct number of significant figures. However, you should round off at the end of all the mathematical operations in order to avoid round-off error. The best way to do calculations is to keep track of the correct number of significant figures during intermediate steps, but round off at the end. For this problem, we underlined the last significant figure in the intermediate steps.

$$\frac{2.526}{3.1} + \frac{0.470}{0.623} + \frac{80.705}{0.4326} = 0.81\underline{48} + 0.75\underline{44} + 186.\underline{558} = 188.1$$

b. Here, the mathematical operation requires that we apply the addition/subtraction rule first, then apply the multiplication/division rule.

$$\frac{6.404 \times 2.91}{18.7 - 17.1} = \frac{6.404 \times 2.91}{1.\underline{6}} = 12$$

- c. $6.071 \times 10^{-5} - 8.2 \times 10^{-6} - 0.521 \times 10^{-4} = 60.71 \times 10^{-6} - 8.2 \times 10^{-6} - 52.1 \times 10^{-6}$
 $= 0.\underline{41} \times 10^{-6} = 4 \times 10^{-7}$

$$d. \frac{3.8 \times 10^{-12} + 4.0 \times 10^{-13}}{4 \times 10^{12} + 6.3 \times 10^{13}} = \frac{38 \times 10^{-13} + 4.0 \times 10^{-13}}{4 \times 10^{12} + 63 \times 10^{12}} = \frac{42 \times 10^{-13}}{67 \times 10^{12}} = 6.3 \times 10^{-26}$$

$$e. \frac{9.5 + 4.1 + 2.8 + 3.175}{4} = \frac{19.575}{4} = 4.89 = 4.9$$

Uncertainty appears in the first decimal place. The average of several numbers can only be as precise as the least precise number. Averages can be exceptions to the significant figure rules.

$$f. \frac{8.925 - 8.905}{8.925} \times 100 = \frac{0.020}{8.925} \times 100 = 0.22$$

$$42. \quad a. 6.022 \times 10^{23} \times 1.05 \times 10^2 = 6.32 \times 10^{25}$$

$$b. \frac{6.6262 \times 10^{-34} \times 2.998 \times 10^8}{2.54 \times 10^{-9}} = 7.82 \times 10^{-17}$$

$$c. 1.285 \times 10^{-2} + 1.24 \times 10^{-3} + 1.879 \times 10^{-1} \\ = 0.1285 \times 10^{-1} + 0.0124 \times 10^{-1} + 1.879 \times 10^{-1} = 2.020 \times 10^{-1}$$

When the exponents are different, it is easiest to apply the addition/subtraction rule when all numbers are based on the same power of 10.

$$d. \frac{(1.00866 - 1.00728)}{6.02205 \times 10^{23}} = \frac{0.00138}{6.02205 \times 10^{23}} = 2.29 \times 10^{-27}$$

$$e. \frac{9.875 \times 10^2 - 9.795 \times 10^2}{9.875 \times 10^2} \times 100 = \frac{0.080 \times 10^2}{9.875 \times 10^2} \times 100 = 8.1 \times 10^{-1}$$

$$f. \frac{9.42 \times 10^2 + 8.234 \times 10^2 + 1.625 \times 10^3}{3} = \frac{0.942 \times 10^3 + 0.824 \times 10^3 + 1.625 \times 10^3}{3} \\ = 1.130 \times 10^3$$

$$43. \quad a. 8.43 \text{ cm} \times \frac{1 \text{ m}}{100 \text{ cm}} \times \frac{1000 \text{ mm}}{\text{m}} = 84.3 \text{ mm} \quad b. 2.41 \times 10^2 \text{ cm} \times \frac{1 \text{ m}}{100 \text{ cm}} = 2.41 \text{ m}$$

$$c. 294.5 \text{ nm} \times \frac{1 \text{ m}}{1 \times 10^9 \text{ nm}} \times \frac{100 \text{ cm}}{\text{m}} = 2.945 \times 10^{-5} \text{ cm}$$

$$d. 1.445 \times 10^4 \text{ m} \times \frac{1 \text{ km}}{1000 \text{ m}} = 14.45 \text{ km} \quad e. 235.3 \text{ m} \times \frac{1000 \text{ mm}}{\text{m}} = 2.353 \times 10^5 \text{ mm}$$

$$f. 903.3 \text{ nm} \times \frac{1 \text{ m}}{1 \times 10^9 \text{ nm}} \times \frac{1 \times 10^6 \mu\text{m}}{\text{m}} = 0.9033 \mu\text{m}$$

$$44. \quad a. 1 \text{ Tg} \times \frac{1 \times 10^{12} \text{ g}}{\text{Tg}} \times \frac{1 \text{ kg}}{1000 \text{ g}} = 1 \times 10^9 \text{ kg}$$

- b. $6.50 \times 10^2 \text{ Tm} \times \frac{1 \times 10^{12} \text{ m}}{\text{Tm}} \times \frac{1 \times 10^9 \text{ nm}}{\text{m}} = 6.50 \times 10^{23} \text{ nm}$
- c. $25 \text{ fg} \times \frac{1 \text{ g}}{1 \times 10^{15} \text{ fg}} \times \frac{1 \text{ kg}}{1000 \text{ g}} = 25 \times 10^{-18} \text{ kg} = 2.5 \times 10^{-17} \text{ kg}$
- d. $8.0 \text{ dm}^3 \times \frac{1 \text{ L}}{\text{dm}^3} = 8.0 \text{ L}$ (1 L = 1 dm³ = 1000 cm³ = 1000 mL)
- e. $1 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{1 \times 10^6 \mu\text{L}}{\text{L}} = 1 \times 10^3 \mu\text{L}$
- f. $1 \mu\text{g} \times \frac{1 \text{ g}}{1 \times 10^6 \mu\text{g}} \times \frac{1 \times 10^{12} \text{ pg}}{\text{g}} = 1 \times 10^6 \text{ pg}$

45. a. Conversion factors are found in Appendix 6. In general, the number of significant figures we use in the conversion factors will be one more than the number of significant figures from the numbers given in the problem. This is usually sufficient to avoid round-off error.

$$3.91 \text{ kg} \times \frac{1 \text{ lb}}{0.4536 \text{ kg}} = 8.62 \text{ lb}; \quad 0.62 \text{ lb} \times \frac{16 \text{ oz}}{\text{lb}} = 9.9 \text{ oz}$$

Baby's weight = 8 lb and 9.9 oz or, to the nearest ounce, 8 lb and 10. oz.

$$51.4 \text{ cm} \times \frac{1 \text{ in}}{2.54 \text{ cm}} = 20.2 \text{ in} \approx 20 \frac{1}{4} \text{ in} = \text{baby's height}$$

- b. $25,000 \text{ mi} \times \frac{1.61 \text{ km}}{\text{mi}} = 4.0 \times 10^4 \text{ km}; \quad 4.0 \times 10^4 \text{ km} \times \frac{1000 \text{ m}}{\text{km}} = 4.0 \times 10^7 \text{ m}$
- c. $V = l \times w \times h = 1.0 \text{ m} \times \left(5.6 \text{ cm} \times \frac{1 \text{ m}}{100 \text{ cm}}\right) \times \left(2.1 \text{ dm} \times \frac{1 \text{ m}}{10 \text{ dm}}\right) = 1.2 \times 10^{-2} \text{ m}^3$

$$1.2 \times 10^{-2} \text{ m}^3 \times \left(\frac{10 \text{ dm}}{\text{m}}\right)^3 \times \frac{1 \text{ L}}{\text{dm}^3} = 12 \text{ L}$$

$$12 \text{ L} \times \frac{1000 \text{ cm}^3}{\text{L}} \times \left(\frac{1 \text{ in}}{2.54 \text{ cm}}\right)^3 = 730 \text{ in}^3; \quad 730 \text{ in}^3 \times \left(\frac{1 \text{ ft}}{12 \text{ in}}\right)^3 = 0.42 \text{ ft}^3$$

46. a. $908 \text{ oz} \times \frac{1 \text{ lb}}{16 \text{ oz}} \times \frac{0.4536 \text{ kg}}{\text{lb}} = 25.7 \text{ kg}$
- b. $12.8 \text{ L} \times \frac{1 \text{ qt}}{0.9463 \text{ L}} \times \frac{1 \text{ gal}}{4 \text{ qt}} = 3.38 \text{ gal}$
- c. $125 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{1 \text{ qt}}{0.9463 \text{ L}} = 0.132 \text{ qt}$

- d. $2.89 \text{ gal} \times \frac{4 \text{ qt}}{1 \text{ gal}} \times \frac{1 \text{ L}}{1.057 \text{ qt}} \times \frac{1000 \text{ mL}}{1 \text{ L}} = 1.09 \times 10^4 \text{ mL}$
- e. $4.48 \text{ lb} \times \frac{453.6 \text{ g}}{1 \text{ lb}} = 2.03 \times 10^3 \text{ g}$
- f. $550 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{1.06 \text{ qt}}{\text{L}} = 0.58 \text{ qt}$
47. a. $1.25 \text{ mi} \times \frac{8 \text{ furlongs}}{\text{mi}} = 10.0 \text{ furlongs}; 10.0 \text{ furlongs} \times \frac{40 \text{ rods}}{\text{furlong}} = 4.00 \times 10^2 \text{ rods}$
- $4.00 \times 10^2 \text{ rods} \times \frac{5.5 \text{ yd}}{\text{rod}} \times \frac{36 \text{ in}}{\text{yd}} \times \frac{2.54 \text{ cm}}{\text{in}} \times \frac{1 \text{ m}}{100 \text{ cm}} = 2.01 \times 10^3 \text{ m}$
- $2.01 \times 10^3 \text{ m} \times \frac{1 \text{ km}}{1000 \text{ m}} = 2.01 \text{ km}$
- b. Let's assume we know this distance to ± 1 yard. First, convert 26 miles to yards.
- $26 \text{ mi} \times \frac{5280 \text{ ft}}{\text{mi}} \times \frac{1 \text{ yd}}{3 \text{ ft}} = 45,760. \text{ yd}$
- $26 \text{ mi} + 385 \text{ yd} = 45,760. \text{ yd} + 385 \text{ yd} = 46,145 \text{ yards}$
- $46,145 \text{ yard} \times \frac{1 \text{ rod}}{5.5 \text{ yd}} = 8390.0 \text{ rods}; 8390.0 \text{ rods} \times \frac{1 \text{ furlong}}{40 \text{ rods}} = 209.75 \text{ furlongs}$
- $46,145 \text{ yard} \times \frac{36 \text{ in}}{\text{yd}} \times \frac{2.54 \text{ cm}}{\text{in}} \times \frac{1 \text{ m}}{100 \text{ cm}} = 42,195 \text{ m}; 42,195 \text{ m} \times \frac{1 \text{ km}}{1000 \text{ m}} = 42.195 \text{ km}$
48. a. $1 \text{ ha} \times \frac{10,000 \text{ m}^2}{\text{ha}} \times \left(\frac{1 \text{ km}}{1000 \text{ m}} \right)^2 = 1 \times 10^{-2} \text{ km}^2$
- b. $5.5 \text{ acre} \times \frac{160 \text{ rod}^2}{\text{acre}} \times \left(\frac{5.5 \text{ yd}}{\text{rod}} \times \frac{36 \text{ in}}{\text{yd}} \times \frac{2.54 \text{ cm}}{\text{in}} \times \frac{1 \text{ m}}{100 \text{ cm}} \right)^2 = 2.2 \times 10^4 \text{ m}^2$
- $2.2 \times 10^4 \text{ m}^2 \times \frac{1 \text{ ha}}{1 \times 10^4 \text{ m}^2} = 2.2 \text{ ha}; 2.2 \times 10^4 \text{ m}^2 \times \left(\frac{1 \text{ km}}{1000 \text{ m}} \right)^2 = 0.022 \text{ km}^2$
- c. Area of lot = $120 \text{ ft} \times 75 \text{ ft} = 9.0 \times 10^3 \text{ ft}^2$
- $9.0 \times 10^3 \text{ ft}^2 \times \left(\frac{1 \text{ yd}}{3 \text{ ft}} \times \frac{1 \text{ rod}}{5.5 \text{ yd}} \right)^2 \times \frac{1 \text{ acre}}{160 \text{ rod}^2} = 0.21 \text{ acre}; \frac{\$6,500}{0.21 \text{ acre}} = \frac{\$31,000}{\text{acre}}$

We can use our result from (b) to get the conversion factor between acres and hectares (5.5 acre = 2.2 ha.). Thus 1 ha = 2.5 acre.

$$0.21 \text{ acre} \times \frac{1 \text{ ha}}{2.5 \text{ acre}} = 0.084 \text{ ha}; \text{ the price is: } \frac{\$6,500}{0.084 \text{ ha}} = \frac{\$77,000}{\text{ha}}$$

$$49. \quad \text{a. } 1 \text{ troy lb} \times \frac{12 \text{ troy oz}}{\text{troy lb}} \times \frac{20 \text{ pw}}{\text{troy oz}} \times \frac{24 \text{ grains}}{\text{pw}} \times \frac{0.0648 \text{ g}}{\text{grain}} \times \frac{1 \text{ kg}}{1000 \text{ g}} = 0.373 \text{ kg}$$

$$1 \text{ troy lb} = 0.373 \text{ kg} \times \frac{2.205 \text{ lb}}{\text{kg}} = 0.822 \text{ lb}$$

$$\text{b. } 1 \text{ troy oz} \times \frac{20 \text{ pw}}{\text{troy oz}} \times \frac{24 \text{ grains}}{\text{pw}} \times \frac{0.0648 \text{ g}}{\text{grain}} = 31.1 \text{ g}$$

$$1 \text{ troy oz} = 31.1 \text{ g} \times \frac{1 \text{ carat}}{0.200 \text{ g}} = 156 \text{ carats}$$

$$\text{c. } 1 \text{ troy lb} = 0.373 \text{ kg}; 0.373 \text{ kg} \times \frac{1000 \text{ g}}{\text{kg}} \times \frac{1 \text{ cm}^3}{19.3 \text{ g}} = 19.3 \text{ cm}^3$$

$$50. \quad \text{a. } 1 \text{ grain ap} \times \frac{1 \text{ scruple}}{20 \text{ grain ap}} \times \frac{1 \text{ dram ap}}{3 \text{ scruples}} \times \frac{3.888 \text{ g}}{\text{dram ap}} = 0.06480 \text{ g}$$

From the previous question, we are given that 1 grain troy = 0.0648 g = 1 grain ap. So the two are the same.

$$\text{b. } 1 \text{ oz ap} \times \frac{8 \text{ dram ap}}{\text{oz ap}} \times \frac{3.888 \text{ g}}{\text{dram ap}} \times \frac{1 \text{ oz troy}^*}{31.1 \text{ g}} = 1.00 \text{ oz troy}; \quad * \text{see Exercise 49b.}$$

$$\text{c. } 5.00 \times 10^2 \text{ mg} \times \frac{1 \text{ g}}{1000 \text{ mg}} \times \frac{1 \text{ dram ap}}{3.888 \text{ g}} \times \frac{3 \text{ scruples}}{\text{dram ap}} = 0.386 \text{ scruple}$$

$$0.386 \text{ scruple} \times \frac{20 \text{ grains ap}}{\text{scruple}} = 7.72 \text{ grains ap}$$

$$\text{d. } 1 \text{ scruple} \times \frac{1 \text{ dram ap}}{3 \text{ scruples}} \times \frac{3.888 \text{ g}}{\text{dram ap}} = 1.296 \text{ g}$$

$$51. \quad 15.6 \text{ g} \times \frac{1 \text{ capsule}}{0.65 \text{ g}} = 24 \text{ capsules}$$

$$52. \quad 1.5 \text{ teaspoons} \times \frac{80. \text{ mg acet}}{0.50 \text{ teaspoon}} = 240 \text{ mg acetaminophen}$$

$$\frac{240 \text{ mg acet}}{24 \text{ lb}} \times \frac{1 \text{ lb}}{0.454 \text{ kg}} = 22 \text{ mg acetaminophen/kg}$$

$$\frac{240 \text{ mg acet}}{35 \text{ lb}} \times \frac{1 \text{ lb}}{0.454 \text{ kg}} = 15 \text{ mg acetaminophen/kg}$$

The range is from 15 to 22 mg acetaminophen per kg of body weight.

$$53. \quad \text{warp } 1.71 = \left(5.00 \times \frac{3.00 \times 10^8 \text{ m}}{\text{s}} \right) \times \frac{1.094 \text{ yd}}{\text{m}} \times \frac{60 \text{ s}}{\text{min}} \times \frac{60 \text{ min}}{\text{h}} \times \frac{1 \text{ knot}}{2030 \text{ yd/h}}$$

$$= 2.91 \times 10^9 \text{ knots}$$

$$\left(5.00 \times \frac{3.00 \times 10^8 \text{ m}}{\text{s}} \right) \times \frac{1 \text{ km}}{1000 \text{ m}} \times \frac{1 \text{ mi}}{1.609 \text{ km}} \times \frac{60 \text{ s}}{\text{min}} \times \frac{60 \text{ min}}{\text{h}} = 3.36 \times 10^9 \text{ mi/h}$$

$$54. \quad \frac{100. \text{ m}}{9.58 \text{ s}} = 10.4 \text{ m/s}; \quad \frac{100. \text{ m}}{9.58 \text{ s}} \times \frac{1 \text{ km}}{1000 \text{ m}} \times \frac{60 \text{ s}}{\text{min}} \times \frac{60 \text{ min}}{\text{h}} = 37.6 \text{ km/h}$$

$$\frac{100. \text{ m}}{9.58 \text{ s}} \times \frac{1.0936 \text{ yd}}{\text{m}} \times \frac{3 \text{ ft}}{\text{yd}} = 34.2 \text{ ft/s}; \quad \frac{34.2 \text{ ft}}{\text{s}} \times \frac{1 \text{ mi}}{5280 \text{ ft}} \times \frac{60 \text{ s}}{\text{min}} \times \frac{60 \text{ min}}{\text{h}} = 23.3 \text{ mi/h}$$

$$1.00 \times 10^2 \text{ yd} \times \frac{1 \text{ m}}{1.0936 \text{ yd}} \times \frac{9.58 \text{ s}}{100. \text{ m}} = 8.76 \text{ s}$$

$$55. \quad 1 \text{ s} \times \frac{1 \text{ min}}{60 \text{ s}} \times \frac{1 \text{ h}}{60 \text{ min}} \times \frac{65 \text{ mi}}{\text{h}} \times \frac{5280 \text{ ft}}{\text{mi}} = 95.3 \text{ ft} = 100 \text{ ft}$$

If you take your eyes off the road for one second traveling at 65 mph, your car travels approximately 100 feet.

$$56. \quad 112 \text{ km} \times \frac{0.6214 \text{ mi}}{\text{km}} \times \frac{1 \text{ h}}{65 \text{ mi}} = 1.1 \text{ h} = 1 \text{ h and } 6 \text{ min}$$

$$112 \text{ km} \times \frac{0.6214 \text{ mi}}{\text{km}} \times \frac{1 \text{ gal}}{28 \text{ mi}} \times \frac{3.785 \text{ L}}{\text{gal}} = 9.4 \text{ L of gasoline}$$

$$57. \quad 180 \text{ lb} \times \frac{1 \text{ kg}}{2.205 \text{ lb}} \times \frac{8.0 \text{ mg}}{\text{kg}} = 650 \text{ mg antibiotic/dose}$$

$$2 \text{ wk} \times \frac{7 \text{ days}}{\text{wk}} \times \frac{2 \text{ doses}}{\text{day}} \times \frac{650 \text{ mg}}{\text{dose}} = 18,000 \text{ mg} = 18 \text{ g antibiotic in total}$$

58. For the gasoline car:

$$500. \text{ mi} \times \frac{1 \text{ gal}}{28.0 \text{ mi}} \times \frac{\$3.50}{\text{gal}} = \$62.5$$

For the E85 car:

$$500. \text{ mi} \times \frac{1 \text{ gal}}{22.5 \text{ mi}} \times \frac{\$2.85}{\text{gal}} = \$63.3$$

The E85 vehicle would cost slightly more to drive 500. miles as compared to the gasoline vehicle (\$63.3 versus \$62.5).

59. Volume of lake = $100 \text{ mi}^2 \times \left(\frac{5280 \text{ ft}}{\text{mi}}\right)^2 \times 20 \text{ ft} = 6 \times 10^{10} \text{ ft}^3$
- $$6 \times 10^{10} \text{ ft}^3 \times \left(\frac{12 \text{ in}}{\text{ft}} \times \frac{2.54 \text{ cm}}{\text{in}}\right)^3 \times \frac{1 \text{ mL}}{\text{cm}^3} \times \frac{0.4 \mu\text{g}}{\text{mL}} = 7 \times 10^{14} \mu\text{g mercury}$$
- $$7 \times 10^{14} \mu\text{g} \times \frac{1 \text{ g}}{1 \times 10^6 \mu\text{g}} \times \frac{1 \text{ kg}}{1 \times 10^3 \text{ g}} = 7 \times 10^5 \text{ kg of mercury}$$
60. Volume of room = $18 \text{ ft} \times 12 \text{ ft} \times 8 \text{ ft} = 1700 \text{ ft}^3$ (carrying one extra significant figure)
- $$1700 \text{ ft}^3 \times \left(\frac{12 \text{ in}}{\text{ft}}\right)^3 \times \left(\frac{2.54 \text{ cm}}{\text{in}}\right)^3 \times \left(\frac{1 \text{ m}}{100 \text{ cm}}\right)^3 = 48 \text{ m}^3$$
- $$48 \text{ m}^3 \times \frac{400,000 \mu\text{g CO}}{\text{m}^3} \times \frac{1 \text{ g CO}}{1 \times 10^6 \mu\text{g CO}} = 19 \text{ g} = 20 \text{ g CO (to 1 sig. fig.)}$$

Temperature

61. a. $T_C = \frac{5}{9} (T_F - 32) = \frac{5}{9} (-459^\circ\text{F} - 32) = -273^\circ\text{C}$; $T_K = T_C + 273 = -273^\circ\text{C} + 273 = 0 \text{ K}$
- b. $T_C = \frac{5}{9} (-40.^\circ\text{F} - 32) = -40.^\circ\text{C}$; $T_K = -40.^\circ\text{C} + 273 = 233 \text{ K}$
- c. $T_C = \frac{5}{9} (68^\circ\text{F} - 32) = 20.^\circ\text{C}$; $T_K = 20.^\circ\text{C} + 273 = 293 \text{ K}$
- d. $T_C = \frac{5}{9} (7 \times 10^7^\circ\text{F} - 32) = 4 \times 10^7^\circ\text{C}$; $T_K = 4 \times 10^7^\circ\text{C} + 273 = 4 \times 10^7 \text{ K}$
62. $96.1^\circ\text{F} \pm 0.2^\circ\text{F}$; first, convert 96.1°F to $^\circ\text{C}$. $T_C = \frac{5}{9} (T_F - 32) = \frac{5}{9} (96.1 - 32) = 35.6^\circ\text{C}$

A change in temperature of 9°F is equal to a change in temperature of 5°C . So the uncertainty is:

$$\pm 0.2^\circ\text{F} \times \frac{5^\circ\text{C}}{9^\circ\text{F}} = \pm 0.1^\circ\text{C}. \text{ Thus } 96.1 \pm 0.2^\circ\text{F} = 35.6 \pm 0.1^\circ\text{C}.$$

63. a. $T_F = \frac{9}{5} \times T_C + 32 = \frac{9}{5} \times 39.2^\circ\text{C} + 32 = 102.6^\circ\text{F}$ (Note: 32 is exact.)
- $$T_K = T_C + 273.2 = 39.2 + 273.2 = 312.4 \text{ K}$$
- b. $T_F = \frac{9}{5} \times (-25) + 32 = -13^\circ\text{F}$; $T_K = -25 + 273 = 248 \text{ K}$

- c. $T_F = \frac{9}{5} \times (-273) + 32 = -459^\circ\text{F}$; $T_K = -273 + 273 = 0 \text{ K}$
- d. $T_F = \frac{9}{5} \times 801 + 32 = 1470^\circ\text{F}$; $T_K = 801 + 273 = 1074 \text{ K}$
64. a. $T_C = T_K - 273 = 233 - 273 = -40.^\circ\text{C}$
 $T_F = \frac{9}{5} \times T_C + 32 = \frac{9}{5} \times (-40.) + 32 = -40.^\circ\text{F}$
- b. $T_C = 4 - 273 = -269^\circ\text{C}$; $T_F = \frac{9}{5} \times (-269) + 32 = -452^\circ\text{F}$
- c. $T_C = 298 - 273 = 25^\circ\text{C}$; $T_F = \frac{9}{5} \times 25 + 32 = 77^\circ\text{F}$
- d. $T_C = 3680 - 273 = 3410^\circ\text{C}$; $T_F = \frac{9}{5} \times 3410 + 32 = 6170^\circ\text{F}$
65. $T_F = \frac{9}{5} \times T_C + 32$; from the problem, we want the temperature where $T_F = 2T_C$.
- Substituting:
- $$2T_C = \frac{9}{5} \times T_C + 32, (0.2)T_C = 32, T_C = \frac{32}{0.2} = 160^\circ\text{C}$$
- $T_F = 2T_C$ when the temperature in Fahrenheit is $2(160) = 320^\circ\text{F}$. Because all numbers when solving the equation are exact numbers, the calculated temperatures are also exact numbers.
66. $T_C = \frac{5}{9}(T_F - 32) = \frac{5}{9}(72 - 32) = 22^\circ\text{C}$
- $T_C = T_K - 273 = 313 - 273 = 40.^\circ\text{C}$
- The difference in temperature between Jupiter at 313 K and Earth at 72°F is $40.^\circ\text{C} - 22^\circ\text{C} = 18^\circ\text{C}$.
67. a. A change in temperature of 140°C is equal to 50°X . Therefore, $\frac{140^\circ\text{C}}{50^\circ\text{X}}$ is the unit conversion between a degree on the X scale to a degree on the Celsius scale. To account for the different zero points, -10° must be subtracted from the temperature on the X scale to get to the Celsius scale. The conversion between $^\circ\text{X}$ to $^\circ\text{C}$ is:
- $$T_C = T_X \times \frac{140^\circ\text{C}}{50^\circ\text{X}} - 10^\circ\text{C}, T_C = T_X \times \frac{14^\circ\text{C}}{5^\circ\text{X}} - 10^\circ\text{C}$$
- The conversion between $^\circ\text{C}$ to $^\circ\text{X}$ would be:
- $$T_X = (T_C + 10^\circ\text{C}) \frac{5^\circ\text{X}}{14^\circ\text{C}}$$

- b. Assuming 10°C and $\frac{5^\circ\text{X}}{14^\circ\text{C}}$ are exact numbers:

$$T_X = (22.0^\circ\text{C} + 10^\circ\text{C}) \frac{5^\circ\text{X}}{14^\circ\text{C}} = 11.4^\circ\text{X}$$


- c. Assuming exact numbers in the temperature conversion formulas:

$$T_C = 58.0^\circ\text{X} \times \frac{14^\circ\text{C}}{5^\circ\text{X}} - 10^\circ\text{C} = 152^\circ\text{C}$$

$$T_K = 152^\circ\text{C} + 273 = 425 \text{ K}$$

$$T_F = \frac{9^\circ\text{F}}{5^\circ\text{C}} \times 152^\circ\text{C} + 32^\circ\text{F} = 306^\circ\text{F}$$

68. a.



A change in temperature of 160°C equals a change in temperature of 100°A .

So $\frac{160^\circ\text{C}}{100^\circ\text{A}}$ is our unit conversion for a degree change in temperature.

At the freezing point: $0^\circ\text{A} = -45^\circ\text{C}$

Combining these two pieces of information:

$$T_A = (T_C + 45^\circ\text{C}) \times \frac{100^\circ\text{A}}{160^\circ\text{C}} = (T_C + 45^\circ\text{C}) \times \frac{5^\circ\text{A}}{8^\circ\text{C}} \text{ or } T_C = T_A \times \frac{8^\circ\text{C}}{5^\circ\text{A}} - 45^\circ\text{C}$$

b. $T_C = (T_F - 32) \times \frac{5}{9}$; $T_C = T_A \times \frac{8}{5} - 45 = (T_F - 32) \times \frac{5}{9}$

$$T_F - 32 = \frac{9}{5} \times \left(T_A \times \frac{8}{5} - 45 \right) = T_A \times \frac{72}{25} - 81, \quad T_F = T_A \times \frac{72^\circ\text{F}}{25^\circ\text{A}} - 49^\circ\text{F}$$

c. $T_C = T_A \times \frac{8}{5} - 45$ and $T_C = T_A$; so $T_C = T_C \times \frac{8}{5} - 45$, $\frac{3T_C}{5} = 45$, $T_C = 75^\circ\text{C} = 75^\circ\text{A}$

d. $T_C = 86^\circ\text{A} \times \frac{8^\circ\text{C}}{5^\circ\text{A}} - 45^\circ\text{C} = 93^\circ\text{C}$; $T_F = 86^\circ\text{A} \times \frac{72^\circ\text{F}}{25^\circ\text{A}} - 49^\circ\text{F} = 199^\circ\text{F} = 2.0 \times 10^2^\circ\text{F}$

e. $T_A = (45^\circ\text{C} + 45^\circ\text{C}) \times \frac{5^\circ\text{A}}{8^\circ\text{C}} = 56^\circ\text{A}$

Density

$$69. \quad \text{Mass} = 350 \text{ lb} \times \frac{453.6 \text{ g}}{\text{lb}} = 1.6 \times 10^5 \text{ g}; \quad V = 1.2 \times 10^4 \text{ in}^3 \times \left(\frac{2.54 \text{ cm}}{\text{in}} \right)^3 = 2.0 \times 10^5 \text{ cm}^3$$

$$\text{Density} = \frac{\text{mass}}{\text{volume}} = \frac{1 \times 10^5 \text{ g}}{2.0 \times 10^5 \text{ cm}^3} = 0.80 \text{ g/cm}^3$$

Because the material has a density less than water, it will float in water.

$$70. \quad \text{Let } d = \text{density}; \quad d_{\text{cube}} = \frac{140.4 \text{ g}}{(3.00 \text{ cm})^3} = 5.20 \text{ g/cm}^3$$

If this is correct to $\pm 1.00\%$ then the density is $5.20 \pm 0.05 \text{ g/cm}^3$

$$V_{\text{sphere}} = (4/3)\pi r^3 = (4/3)\pi(1.42 \text{ cm})^3 = 12.0 \text{ cm}^3$$

$$d_{\text{sphere}} = \frac{61.6 \text{ g}}{12.0 \text{ cm}^3} = 5.13 \text{ g/cm}^3 = 5.13 \pm 0.05 \text{ g/cm}^3$$

Since d_{cube} is between 5.15 and 5.25 g/cm^3 and d_{sphere} is between 5.08 and 5.18 g/cm^3 , the error limits overlap and we can't decisively determine if they are built of the same material. The data are not precise enough to determine.

$$71. \quad V = \frac{4}{3}\pi r^3 = \frac{4}{3} \times 3.14 \times \left(7.0 \times 10^5 \text{ km} \times \frac{1000 \text{ m}}{\text{km}} \times \frac{100 \text{ cm}}{\text{m}} \right)^3 = 1.4 \times 10^{33} \text{ cm}^3$$

$$\text{Density} = \frac{\text{mass}}{\text{volume}} = \frac{2 \times 10^{36} \text{ kg} \times \frac{1000 \text{ g}}{\text{kg}}}{1.4 \times 10^{33} \text{ cm}^3} = 1.4 \times 10^6 \text{ g/cm}^3 = 1 \times 10^6 \text{ g/cm}^3$$

$$72. \quad V = l \times w \times h = 2.9 \text{ cm} \times 3.5 \text{ cm} \times 10.0 \text{ cm} = 1.0 \times 10^2 \text{ cm}^3$$

$$d = \text{density} = \frac{615.0 \text{ g}}{1.0 \times 10^2 \text{ cm}^3} = \frac{6.2 \text{ g}}{\text{cm}^3}$$

$$73. \quad \text{a.} \quad 5.0 \text{ carat} \times \frac{0.200 \text{ g}}{\text{carat}} \times \frac{1 \text{ cm}^3}{3.51 \text{ g}} = 0.28 \text{ cm}^3$$

$$\text{b.} \quad 2.8 \text{ mL} \times \frac{1 \text{ cm}^3}{\text{mL}} \times \frac{3.51 \text{ g}}{\text{cm}^3} \times \frac{1 \text{ carat}}{0.200 \text{ g}} = 49 \text{ carats}$$

$$74. \quad 1 \text{ mL} = 1 \text{ cm}^3; \quad 125 \text{ cm}^3 \times \frac{3.12 \text{ g}}{\text{cm}^3} = 390. \text{ g Br}_2; \quad 85.0 \text{ g} \times \frac{1 \text{ mL}}{3.12 \text{ g}} = 27.2 \text{ mL Br}_2$$

$$75. \quad V = 21.6 \text{ mL} - 12.7 \text{ mL} = 8.9 \text{ mL}; \quad \text{density} = \frac{33.42 \text{ g}}{8.9 \text{ mL}} = 3.8 \text{ g/mL} = 3.8 \text{ g/cm}^3$$

$$76. \quad 5.25 \text{ g} \times \frac{1 \text{ cm}^3}{10.5 \text{ g}} = 0.500 \text{ cm}^3 = 0.500 \text{ mL}$$

The volume in the cylinder will rise to 11.7 mL (11.2 mL + 0.500 mL = 11.7 mL).

77. a. Both have the same mass of 1.0 kg.

b. 1.0 mL of mercury; mercury is more dense than water. *Note:* 1 mL = 1 cm³.

$$1.0 \text{ mL} \times \frac{13.6 \text{ g}}{\text{mL}} = 14 \text{ g of mercury}; \quad 1.0 \text{ mL} \times \frac{0.998 \text{ g}}{\text{mL}} = 1.0 \text{ g of water}$$

c. Same; both represent 19.3 g of substance.

$$19.3 \text{ mL} \times \frac{0.9982 \text{ g}}{\text{mL}} = 19.3 \text{ g of water}; \quad 1.00 \text{ mL} \times \frac{19.32 \text{ g}}{\text{mL}} = 19.3 \text{ g of gold}$$

d. 1.0 L of benzene (880 g versus 670 g)

$$75 \text{ mL} \times \frac{8.96 \text{ g}}{\text{mL}} = 670 \text{ g of copper}; \quad 1.0 \text{ L} \times \frac{1000 \text{ mL}}{\text{L}} \times \frac{0.880 \text{ g}}{\text{mL}} = 880 \text{ g of benzene}$$

$$78. \quad \text{a.} \quad 1.50 \text{ qt} \times \frac{1 \text{ L}}{1.0567 \text{ qt}} \times \frac{1000 \text{ mL}}{\text{L}} \times \frac{0.789 \text{ g}}{\text{mL}} = 1120 \text{ g ethanol}$$

$$\text{b.} \quad 3.5 \text{ in}^3 \times \left(\frac{2.54 \text{ cm}}{\text{in}} \right)^3 \times \frac{13.6 \text{ g}}{\text{cm}^3} = 780 \text{ g mercury}$$

79. a. 1.0 kg feather; feathers are less dense than lead.

b. 100 g water; water is less dense than gold.

c. Same; both volumes are 1.0 L.

$$80. \quad \text{a.} \quad \text{H}_2(\text{g}): \quad V = 25.0 \text{ g} \times \frac{1 \text{ cm}^3}{0.000084 \text{ g}} = 3.0 \times 10^5 \text{ cm}^3 \quad [\text{H}_2(\text{g}) = \text{hydrogen gas.}]$$

$$\text{b.} \quad \text{H}_2\text{O}(\text{l}): \quad V = 25.0 \text{ g} \times \frac{1 \text{ cm}^3}{0.9982 \text{ g}} = 25.0 \text{ cm}^3 \quad [\text{H}_2\text{O}(\text{l}) = \text{water.}]$$

$$\text{c.} \quad \text{Fe}(\text{s}): \quad V = 25.0 \text{ g} \times \frac{1 \text{ cm}^3}{7.87 \text{ g}} = 3.18 \text{ cm}^3 \quad [\text{Fe}(\text{s}) = \text{iron.}]$$

Notice the huge volume of the gaseous H₂ sample as compared to the liquid and solid samples. The same mass of gas occupies a volume that is over 10,000 times larger than the liquid sample. Gases are indeed mostly empty space.

$$81. \quad V = 1.00 \times 10^3 \text{ g} \times \frac{1 \text{ cm}^3}{22.57 \text{ g}} = 44.3 \text{ cm}^3$$

$$44.3 \text{ cm}^3 = 1 \times w \times h = 4.00 \text{ cm} \times 4.00 \text{ cm} \times h, \quad h = 2.77 \text{ cm}$$

82. $V = 22 \text{ g} \times \frac{1 \text{ cm}^3}{8.96 \text{ g}} = 2.5 \text{ cm}^3$; $V = \pi r^2 \times l$, where l = length of the wire

$$2.5 \text{ cm}^3 = \pi \times \left(\frac{0.25 \text{ mm}}{2}\right)^2 \times \left(\frac{1 \text{ cm}}{10 \text{ mm}}\right)^2 \times l, \quad l = 5.1 \times 10^3 \text{ cm} = 170 \text{ ft}$$

Classification and Separation of Matter

83. A gas has molecules that are very far apart from each other, whereas a solid or liquid has molecules that are very close together. An element has the same type of atom, whereas a compound contains two or more different elements. Picture i represents an element that exists as two atoms bonded together (like H_2 or O_2 or N_2). Picture iv represents a compound (like CO , NO , or HF). Pictures iii and iv contain representations of elements that exist as individual atoms (like Ar, Ne, or He).

- Picture iv represents a gaseous compound. Note that pictures ii and iii also contain a gaseous compound, but they also both have a gaseous element present.
- Picture vi represents a mixture of two gaseous elements.
- Picture v represents a solid element.
- Pictures ii and iii both represent a mixture of a gaseous element and a gaseous compound.

84. Solid: rigid; has a fixed volume and shape; slightly compressible

Liquid: definite volume but no specific shape; assumes shape of the container; slightly compressible

Gas: no fixed volume or shape; easily compressible

Pure substance: has constant composition; can be composed of either compounds or elements

Element: substances that cannot be decomposed into simpler substances by chemical or physical means.

Compound: a substance that can be broken down into simpler substances (elements) by chemical processes.

Homogeneous mixture: a mixture of pure substances that has visibly indistinguishable parts.

Heterogeneous mixture: a mixture of pure substances that has visibly distinguishable parts.

Solution: a homogeneous mixture; can be a solid, liquid or gas

Chemical change: a given substance becomes a new substance or substances with different properties and different composition.

Physical change: changes the form (g, l, or s) of a substance but does not change the chemical composition of the substance.

85. Homogeneous: Having visibly indistinguishable parts (the same throughout).

Heterogeneous: Having visibly distinguishable parts (not uniform throughout).

a. heterogeneous (due to hinges, handles, locks, etc.)

b. homogeneous (hopefully; if you live in a heavily polluted area, air may be heterogeneous.)

c. homogeneous

d. homogeneous (hopefully, if not polluted)

e. heterogeneous

f. heterogeneous

86. a. heterogeneous

b. homogeneous

c. heterogeneous

d. homogeneous (assuming no imperfections in the glass)

e. heterogeneous (has visibly distinguishable parts)

87. a. pure b. mixture c. mixture d. pure e. mixture (copper and zinc)

f. pure g. mixture h. mixture i. mixture

Iron and uranium are elements. Water (H_2O) is a compound because it is made up of two or more different elements. Table salt is usually a homogeneous mixture composed mostly of sodium chloride (NaCl), but will usually contain other substances that help absorb water vapor (an anticaking agent).

88. Initially, a mixture is present. The magnesium and sulfur have only been placed together in the same container at this point, but no reaction has occurred. When heated, a reaction occurs. Assuming the magnesium and sulfur had been measured out in exactly the correct ratio for complete reaction, the remains after heating would be a pure compound composed of magnesium and sulfur. However, if there were an excess of either magnesium or sulfur, the remains after reaction would be a mixture of the compound produced and the excess reactant.

89. Chalk is a compound because it loses mass when heated and appears to change into another substance with different physical properties (the hard chalk turns into a crumbly substance).

90. Because vaporized water is still the *same substance* as solid water (H_2O), no chemical reaction has occurred. Sublimation is a physical change.

91. A physical change is a change in the state of a substance (solid, liquid, and gas are the three states of matter); a physical change does not change the chemical composition of the

substance. A chemical change is a change in which a given substance is converted into another substance having a different formula (composition).

- a. Vaporization refers to a liquid converting to a gas, so this is a physical change. The formula (composition) of the moth ball does not change.
 - b. This is a chemical change since hydrofluoric acid (HF) is reacting with glass (SiO₂) to form new compounds that wash away.
 - c. This is a physical change because all that is happening during the boiling process is the conversion of liquid alcohol to gaseous alcohol. The alcohol formula (C₂H₅OH) does not change.
 - d. This is a chemical change since the acid is reacting with cotton to form new compounds.
- 92.
- a. Distillation separates components of a mixture, so the orange liquid is a mixture (has an average color of the yellow liquid and the red solid). Distillation utilizes boiling point differences to separate out the components of a mixture. Distillation is a physical change because the components of the mixture do not become different compounds or elements.
 - b. Decomposition is a type of chemical reaction. The crystalline solid is a compound, and decomposition is a chemical change where new substances are formed.
 - c. Tea is a mixture of tea compounds dissolved in water. The process of mixing sugar into tea is a physical change. Sugar doesn't react with the tea compounds, it just makes the solution sweeter.

Additional Exercises

93. The object that sinks has a greater density than water and the object that floats has a smaller density than water. Since both objects have the same mass, the sphere that sinks must have the smaller volume which makes it more dense. Therefore, the object that floats has the larger volume along with the greater diameter.

94.
$$422 \text{ mg caffeine} \times \frac{1 \text{ g}}{1000 \text{ mg}} \times \frac{6.02 \times 10^{23} \text{ molecules}}{194 \text{ g caffeine}} = 1.31 \times 10^{21} \text{ caffeine molecules}$$

95. Because each pill is 4.0% Lipitor by mass, for every 100.0 g of pills, there are 4.0 g of Lipitor present. Note that 100 pills is assumed to be an exact number.

$$100 \text{ pills} \times \frac{2.5 \text{ g}}{\text{pill}} \times \frac{4.0 \text{ g Lipitor}}{100.0 \text{ g pills}} \times \frac{1 \text{ kg}}{1000 \text{ g}} = 0.010 \text{ kg Lipitor}$$

96.
$$126 \text{ gal} \times \frac{4 \text{ qt}}{\text{gal}} \times \frac{1 \text{ L}}{1.057 \text{ qt}} = 477 \text{ L}$$

$$97. \quad \text{Total volume} = \left(200. \text{ m} \times \frac{100 \text{ cm}}{\text{m}}\right) \times \left(300. \text{ m} \times \frac{100 \text{ cm}}{\text{m}}\right) \times 4.0 \text{ cm} = 2.4 \times 10^9 \text{ cm}^3$$

Volume of topsoil covered by 1 bag =

$$\left[10. \text{ ft}^2 \times \left(\frac{12 \text{ in}}{\text{ft}}\right)^2 \times \left(\frac{2.54 \text{ cm}}{\text{in}}\right)^2\right] \times \left(1.0 \text{ in} \times \frac{2.54 \text{ cm}}{\text{in}}\right) = 2.4 \times 10^4 \text{ cm}^3$$

$$2.4 \times 10^9 \text{ cm}^3 \times \frac{1 \text{ bag}}{2.4 \times 10^4 \text{ cm}^3} = 1.0 \times 10^5 \text{ bags topsoil}$$

98. a. No; if the volumes were the same, then the gold idol would have a much greater mass because gold is much more dense than sand.

$$b. \quad \text{Mass} = 1.0 \text{ L} \times \frac{1000 \text{ cm}^3}{\text{L}} \times \frac{19.32 \text{ g}}{\text{cm}^3} \times \frac{1 \text{ kg}}{1000 \text{ g}} = 19.32 \text{ kg} (= 42.59 \text{ lb})$$

It wouldn't be easy to play catch with the idol because it would have a mass of over 40 pounds.

$$99. \quad 1 \text{ light year} = 1 \text{ yr} \times \frac{365 \text{ day}}{\text{yr}} \times \frac{24 \text{ h}}{\text{day}} \times \frac{60 \text{ min}}{\text{h}} \times \frac{60 \text{ s}}{\text{min}} \times \frac{186,000 \text{ mi}}{\text{s}} = 5.87 \times 10^{12} \text{ miles}$$

$$9.6 \text{ parsecs} \times \frac{3.26 \text{ light yr}}{\text{parsec}} \times \frac{5.87 \times 10^{12} \text{ mi}}{\text{light yr}} \times \frac{1.609 \text{ km}}{\text{mi}} \times \frac{1000 \text{ m}}{\text{km}} = 3.0 \times 10^{17} \text{ m}$$

$$100. \quad \text{Density} = \frac{\text{mass}}{\text{volume}} = \frac{0.384 \text{ g}}{0.32 \text{ cm}^3} = 1.2 \text{ g/cm}^3; \text{ from the table, the other ingredient is caffeine.}$$

$$101. \quad a. \quad 0.25 \text{ lb} \times \frac{453.6 \text{ g}}{\text{lb}} \times \frac{1.0 \text{ g tryptophan}}{100.0 \text{ g turkey}} = 1.1 \text{ g tryptophan}$$

$$b. \quad 0.25 \text{ qt} \times \frac{0.9463 \text{ L}}{\text{qt}} \times \frac{1.04 \text{ kg}}{\text{L}} \times \frac{1000 \text{ g}}{\text{kg}} \times \frac{2.0 \text{ g tryptophan}}{100.0 \text{ g milk}} = 4.9 \text{ g tryptophan}$$

102. A chemical change involves the change of one or more substances into other substances through a reorganization of the atoms. A physical change involves the change in the form of a substance, but not its chemical composition.

a. physical change (Just smaller pieces of the same substance.)

b. chemical change (Chemical reactions occur.)

c. chemical change (Bonds are broken.)

d. chemical change (Bonds are broken.)

e. physical change (Water is changed from a liquid to a gas.)

f. physical change (Chemical composition does not change.)

$$103. \quad 5.4 \text{ L blood} \times \frac{1000 \text{ mL}}{\text{L}} \times \frac{250 \text{ mg cholesterol}}{100.0 \text{ mL blood}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 13.5 \text{ g} = 14 \text{ g cholesterol}$$

$$104. \quad \text{a. For } \frac{103 \pm 1}{101 \pm 1}: \text{ maximum} = \frac{104}{100} = 1.04; \text{ minimum} = \frac{102}{102} = 1.00$$

$$\text{So: } \frac{103 \pm 1}{101 \pm 1} = 1.02 \pm 0.02$$

$$\text{b. For } \frac{101 \pm 1}{99 \pm 1}: \text{ maximum} = \frac{102}{98} = 1.04; \text{ minimum} = \frac{100}{100} = 1.00$$

$$\text{So: } \frac{101 \pm 1}{99 \pm 1} = 1.02 \pm 0.02$$

$$\text{c. For } \frac{99 \pm 1}{101 \pm 1}: \text{ maximum} = \frac{100}{100} = 1.00; \text{ minimum} = \frac{98}{102} = 0.96$$

$$\text{So: } \frac{99 \pm 1}{101 \pm 1} = 0.98 \pm 0.02$$

Considering the error limits, answers to parts a and b should be expressed to three significant figures while the part c answer should be expressed to two significant figures. Using the multiplication/division rule leads to a different result in part b; according to the rule, the part b answer should be to two significant figures. If this is the case, then the answer to part b is 1.0, which implies the answer to the calculation is somewhere between 0.95 and 1.05. The actual error limit to the answer is better than this, so we should use the more precise way of expressing the answer. The significant figure rules give general guidelines for estimating uncertainty; there are exceptions to the rules.

$$105. \quad 18.5 \text{ cm} \times \frac{10.0^\circ \text{F}}{5.25 \text{ cm}} = 35.2^\circ \text{F increase}; T_{\text{final}} = 98.6 + 35.2 = 133.8^\circ \text{F}$$

$$T_{\text{C}} = 5/9 (133.8 - 32) = 56.56^\circ \text{C}$$

$$106. \quad \text{Mass}_{\text{benzene}} = 58.80 \text{ g} - 25.00 \text{ g} = 33.80 \text{ g}; V_{\text{benzene}} = 33.80 \text{ g} \times \frac{1 \text{ cm}^3}{0.880 \text{ g}} = 38.4 \text{ cm}^3$$

$$V_{\text{solid}} = 50.0 \text{ cm}^3 - 38.4 \text{ cm}^3 = 11.6 \text{ cm}^3; \text{ density} = \frac{25.00 \text{ g}}{11.6 \text{ cm}^3} = 2.16 \text{ g/cm}^3$$

107. a. Volume \times density = mass; the orange block is more dense. Because mass (orange) > mass (blue) and because volume (orange) < volume (blue), the density of the orange block must be greater to account for the larger mass of the orange block.

- b. Which block is more dense cannot be determined. Because mass (orange) > mass (blue) and because volume (orange) > volume (blue), the density of the orange block may or may not be larger than the blue block. If the blue block is more dense, its density cannot be so large that its mass is larger than the orange block's mass.
- c. The blue block is more dense. Because mass (blue) = mass (orange) and because volume (blue) < volume (orange), the density of the blue block must be larger in order to equate the masses.
- d. The blue block is more dense. Because mass (blue) > mass (orange) and because the volumes are equal, the density of the blue block must be larger in order to give the blue block the larger mass.

$$108. \text{ Circumference} = c = 2\pi r; \quad V = \frac{4\pi r^3}{3} = \frac{4\pi}{3} \left(\frac{c}{2\pi} \right)^3 = \frac{c^3}{6\pi^2}$$

$$\text{Largest density} = \frac{5.25 \text{ oz}}{\frac{(9.00 \text{ in})^3}{6\pi^2}} = \frac{5.25 \text{ oz}}{12.3 \text{ in}^3} = \frac{0.427 \text{ oz}}{\text{in}^3}$$

$$\text{Smallest density} = \frac{5.00 \text{ oz}}{\frac{(9.25 \text{ in})^3}{6\pi^2}} = \frac{5.00 \text{ oz}}{13.4 \text{ in}^3} = \frac{0.373 \text{ oz}}{\text{in}^3}$$

$$\text{Maximum range is: } \frac{(0.373 - 0.427) \text{ oz}}{\text{in}^3} \text{ or } 0.40 \pm 0.03 \text{ oz/in}^3$$

Uncertainty is in 2nd decimal place.

$$109. \quad V = V_{\text{final}} - V_{\text{initial}}; \quad d = \frac{28.90 \text{ g}}{9.8 \text{ cm}^3 - 6.4 \text{ cm}^3} = \frac{28.90 \text{ g}}{3.4 \text{ cm}^3} = 8.5 \text{ g/cm}^3$$

$$d_{\text{max}} = \frac{\text{mass}_{\text{max}}}{V_{\text{min}}}; \quad \text{we get } V_{\text{min}} \text{ from } 9.7 \text{ cm}^3 - 6.5 \text{ cm}^3 = 3.2 \text{ cm}^3.$$

$$d_{\text{max}} = \frac{28.93 \text{ g}}{3.2 \text{ cm}^3} = \frac{9.0 \text{ g}}{\text{cm}^3}; \quad d_{\text{min}} = \frac{\text{mass}_{\text{min}}}{V_{\text{max}}} = \frac{28.87 \text{ g}}{9.9 \text{ cm}^3 - 6.3 \text{ cm}^3} = \frac{8.0 \text{ g}}{\text{cm}^3}$$

The density is $8.5 \pm 0.5 \text{ g/cm}^3$.

110. We need to calculate the maximum and minimum values of the density, given the uncertainty in each measurement. The maximum value is:

$$d_{\text{max}} = \frac{19.625 \text{ g} + 0.002 \text{ g}}{25.00 \text{ cm}^3 - 0.03 \text{ cm}^3} = \frac{19.627 \text{ g}}{24.97 \text{ cm}^3} = 0.7860 \text{ g/cm}^3$$

The minimum value of the density is:

$$d_{\min} = \frac{19.625 \text{ g} - 0.002 \text{ g}}{25.00 \text{ cm}^3 + 0.03 \text{ cm}^3} = \frac{19.623 \text{ g}}{25.03 \text{ cm}^3} = 0.7840 \text{ g/cm}^3$$

The density of the liquid is between 0.7840 and 0.7860 g/cm³. These measurements are sufficiently precise to distinguish between ethanol (d = 0.789 g/cm³) and isopropyl alcohol (d = 0.785 g/cm³).

ChemWork Problems

$$111. \quad 4145 \text{ mi} \times \frac{5280 \text{ ft}}{\text{mi}} \times \frac{1 \text{ fathom}}{6 \text{ ft}} \times \frac{1 \text{ cable length}}{100 \text{ fathom}} = 3.648 \times 10^4 \text{ cable lengths}$$

$$4145 \text{ mi} \times \frac{1 \text{ km}}{0.62137 \text{ mi}} \times \frac{1000 \text{ m}}{\text{km}} = 6.671 \times 10^6 \text{ m}$$

$$3.648 \times 10^4 \text{ cable lengths} \times \frac{1 \text{ nautical mile}}{10 \text{ cable lengths}} = 3,648 \text{ nautical miles}$$

$$112. \quad \frac{1.25 \text{ mi}}{119.2 \text{ s}} \times \frac{1 \text{ km}}{0.6214 \text{ mi}} \times \frac{1000 \text{ m}}{1 \text{ km}} = 16.9 \text{ m/s}$$

$$113. \quad T_{\text{C}} = \frac{5}{9}(T_{\text{F}} - 32) = \frac{5}{9}(134^{\circ}\text{F} - 32) = 56.7^{\circ}\text{C}; \text{ phosphorus would be a liquid.}$$




$$114. \quad V = \frac{4}{3}\pi r^3 = \frac{4}{3} \times 3.14 \times \left(69 \text{ pm} \times \frac{1 \times 10^{-12} \text{ m}}{\text{pm}} \times \frac{100 \text{ cm}}{\text{m}} \right)^3 = 1.4 \times 10^{-24} \text{ cm}^3$$

$$\text{Density} = \frac{\text{mass}}{\text{volume}} = \frac{3.35 \times 10^{-23} \text{ g}}{1.4 \times 10^{-24} \text{ cm}^3} = 24 \text{ g/cm}^3$$

115. a. False; sugar is generally considered to be the pure compound sucrose, C₁₂H₂₂O₁₁.
 b. False; elements and compounds are pure substances.
 c. True; air is a mixture of mostly nitrogen and oxygen gases.
 d. False; gasoline has many additives, so it is a mixture.
 e. True; compounds are broken down to elements by chemical change.
116. The rusting of iron is the only change listed where chemical formulas change, so it is the only chemical change. The others are physical properties. Note that the red glow of a platinum wire assumes no reaction between platinum and oxygen; the red glow is just hot Pt.

Challenge Problems

117. In a subtraction, the result gets smaller, but the uncertainties add. If the two numbers are very close together, the uncertainty may be larger than the result. For example, let's assume we want to take the difference of the following two measured quantities, $999,999 \pm 2$ and $999,996 \pm 2$. The difference is 3 ± 4 . Because of the uncertainty, subtracting two similar numbers is poor practice.
118. In general, glassware is estimated to one place past the markings.

<p>a. 128.7 mL glassware</p>  <p style="text-align: center;">read to tenth's place</p>	<p>b. 18 mL glassware</p>  <p style="text-align: center;">read to one's place</p>	<p>c. 23.45 mL glassware</p>  <p style="text-align: center;">read to two decimal places</p>
---	--	--

Total volume = $128.7 + 18 + 23.45 = 170.15 = 170$. (Due to 18, the sum would be known only to the ones place.)

119. a. $\frac{2.70 - 2.64}{2.70} \times 100 = 2\%$ b. $\frac{|16.12 - 16.48|}{16.12} \times 100 = 2.2\%$
- c. $\frac{1.000 - 0.9981}{1.000} \times 100 = \frac{0.002}{1.000} \times 100 = 0.2\%$
120. a. At some point in 1982, the composition of the metal used in minting pennies was changed because the mass changed during this year (assuming the volume of the pennies were constant).
- b. It should be expressed as 3.08 ± 0.05 g. The uncertainty in the second decimal place will swamp any effect of the next decimal places.
121. Heavy pennies (old): mean mass = 3.08 ± 0.05 g

$$\text{Light pennies (new): mean mass} = \frac{(2.467 + 2.545 + 2.518)}{3} = 2.51 \pm 0.04 \text{ g}$$

Because we are assuming that volume is additive, let's calculate the volume of 100.0 g of each type of penny, then calculate the density of the alloy. For 100.0 g of the old pennies, 95 g will be Cu (copper) and 5 g will be Zn (zinc).

$$V = 95 \text{ g Cu} \times \frac{1 \text{ cm}^3}{8.96 \text{ g}} + 5 \text{ g Zn} \times \frac{1 \text{ cm}^3}{7.14 \text{ g}} = 11.3 \text{ cm}^3 \text{ (carrying one extra sig. fig.)}$$

$$\text{Density of old pennies} = \frac{100. \text{ g}}{11.3 \text{ cm}^3} = 8.8 \text{ g/cm}^3$$

For 100.0 g of new pennies, 97.6 g will be Zn and 2.4 g will be Cu.

$$V = 2.4 \text{ g Cu} \times \frac{1 \text{ cm}^3}{8.96 \text{ g}} + 97.6 \text{ g Zn} \times \frac{1 \text{ cm}^3}{7.14 \text{ g}} = 13.94 \text{ cm}^3 \text{ (carrying one extra sig. fig.)}$$

$$\text{Density of new pennies} = \frac{100. \text{ g}}{13.94 \text{ cm}^3} = 7.17 \text{ g/cm}^3$$

$d = \frac{\text{mass}}{\text{volume}}$; because the volume of both types of pennies are assumed equal, then:

$$\frac{d_{\text{new}}}{d_{\text{old}}} = \frac{\text{mass}_{\text{new}}}{\text{mass}_{\text{old}}} = \frac{7.17 \text{ g/cm}^3}{8.8 \text{ g/cm}^3} = 0.81$$

The calculated average mass ratio is: $\frac{\text{mass}_{\text{new}}}{\text{mass}_{\text{old}}} = \frac{2.51 \text{ g}}{3.08 \text{ g}} = 0.815$

To the first two decimal places, the ratios are the same. If the assumptions are correct, then we can reasonably conclude that the difference in mass is accounted for by the difference in alloy used.


122. a. At 8 a.m., approximately 57 cars pass through the intersection per hour.
- b. At 12 a.m. (midnight), only 1 or 2 cars pass through the intersection per hour.
- c. Traffic at the intersection is limited to less than 10 cars per hour from 8 p.m. to 5 a.m. Starting at 6 a.m., there is a steady increase in traffic through the intersection, peaking at 8 a.m. when approximately 57 cars pass per hour. Past 8 a.m. traffic moderates to about 40 cars through the intersection per hour until noon, and then decreases to 21 cars per hour by 3 p.m. Past 3 p.m. traffic steadily increases to a peak of 52 cars per hour at 5 p.m., and then steadily decreases to the overnight level of less than 10 cars through the intersection per hour.
- d. The traffic pattern through the intersection is directly related to the work schedules of the general population as well as to the store hours of the businesses in downtown.
- e. Run the same experiment on a Sunday, when most of the general population doesn't work and when a significant number of downtown stores are closed in the morning.
123. Let x = mass of copper and y = mass of silver.

$$105.0 \text{ g} = x + y \text{ and } 10.12 \text{ mL} = \frac{x}{8.96} + \frac{y}{10.5}; \text{ solving and carrying 1 extra sig. fig.:}$$


$$\left(10.12 = \frac{x}{8.96} + \frac{105.0 - x}{10.5} \right) \times 8.96 \times 10.5, \quad 952.1 = (10.5)x + 940.8 - (8.96)x$$

$$11.3 = (1.54)x, \quad x = 7.3 \text{ g}; \quad \text{mass \% Cu} = \frac{7.3 \text{ g}}{105.0 \text{ g}} \times 100 = 7.0\% \text{ Cu}$$

124. a.



b.



125. a. One possibility is that rope B is not attached to anything and rope A and rope C are connected via a pair of pulleys and/or gears.
- b. Try to pull rope B out of the box. Measure the distance moved by C for a given movement of A. Hold either A or C firmly while pulling on the other rope.
126. The bubbles of gas is air in the sand that is escaping; methanol and sand are not reacting. We will assume that the mass of trapped air is insignificant.

$$\text{Mass of dry sand} = 37.3488 \text{ g} - 22.8317 \text{ g} = 14.5171 \text{ g}$$

$$\text{Mass of methanol} = 45.2613 \text{ g} - 37.3488 \text{ g} = 7.9125 \text{ g}$$

$$\text{Volume of sand particles (air absent)} = \text{volume of sand and methanol} - \text{volume of methanol}$$

$$\text{Volume of sand particles (air absent)} = 17.6 \text{ mL} - 10.00 \text{ mL} = 7.6 \text{ mL}$$

$$\text{Density of dry sand (air present)} = \frac{14.5171 \text{ g}}{10.0 \text{ mL}} = 1.45 \text{ g/mL}$$

$$\text{Density of methanol} = \frac{7.9125 \text{ g}}{10.00 \text{ mL}} = 0.7913 \text{ g/mL}$$

$$\text{Density of sand particles (air absent)} = \frac{14.5171 \text{ g}}{7.6 \text{ mL}} = 1.9 \text{ g/mL}$$

Integrative Problems

127. 2.97×10^8 persons \times 0.0100 = 2.97×10^6 persons contributing

$$\frac{\$4.75 \times 10^8}{2.97 \times 10^6 \text{ persons}} = \$160./\text{person}; \frac{\$160.}{\text{person}} \times \frac{20 \text{ nickels}}{\$1} = 3.20 \times 10^3 \text{ nickels/person}$$

$$\frac{\$160.}{\text{person}} \times \frac{1 \text{ £}}{\$1.869} = 85.6 \text{ £/person}$$

128. $\frac{22,610 \text{ kg}}{\text{m}^3} \times \frac{1000 \text{ g}}{\text{kg}} \times \frac{1 \text{ m}^3}{1 \times 10^6 \text{ cm}^3} = 22.61 \text{ g/cm}^3$

$$\text{Volume of block} = 10.0 \text{ cm} \times 8.0 \text{ cm} \times 9.0 \text{ cm} = 720 \text{ cm}^3; \frac{22.61 \text{ g}}{\text{cm}^3} \times 720 \text{ cm}^3 = 1.6 \times 10^4 \text{ g}$$

129. At 200.0°F: $T_C = \frac{5}{9}(200.0^\circ\text{F} - 32^\circ\text{F}) = 93.33^\circ\text{C}$; $T_K = 93.33 + 273.15 = 366.48 \text{ K}$

$$\text{At } -100.0^\circ\text{F: } T_C = \frac{5}{9}(-100.0^\circ\text{F} - 32^\circ\text{F}) = -73.33^\circ\text{C}; T_K = -73.33^\circ\text{C} + 273.15 = 199.82 \text{ K}$$

$$\Delta T(^{\circ}\text{C}) = [93.33^\circ\text{C} - (-73.33^\circ\text{C})] = 166.66^\circ\text{C}; \Delta T(\text{K}) = (366.48 \text{ K} - 199.82 \text{ K}) = 166.66 \text{ K}$$

The “300 Club” name only works for the Fahrenheit scale; it does not hold true for the Celsius and Kelvin scales.